

EFFICIENCY OF THE PHRAGMITES AUSTRALIS AND TYPHA DOMINGENSIS ROOTS IN REMEDIATION OF POLYCYCLIC AROMATIC HYDROCARBONS (PAHS) FROM FRESHWATER SEDIMENTS

S. W. Alwan

Lecturer

College of Agriculture/ Al-Qadissyah University

abo_hussian@yahoo.com

ABSTRACT

This study had been conducted to quantify the concentrations of PAHs compounds in the roots of *Phragmites australis* [Cav.] Trin. ex Steudel) and *Typha domingensis* Pers. and to assess their efficiency in remediation of these pollutants from the sediment of fresh water sediment in Hilla, Daghara and Diwaniyah River. Samples (whole plants) were collected from the selective sites, seasonally during 2011-2012. Results showed the efficiency of *Phragmites australis* and *Typha domingensis* roots in the accumulation of HMW-PAHs, which were recorded (69.9 and 35.67 ng.g⁻¹ dry weight DW) during winter and spring whereas these values were (21.8 and 38.7 ng.g⁻¹) in shoots samples during the summer and autumn for both plants respectively. In contrast, LMW-PAHs compounds were not detected in some seasons and sites whereas they were recorded (4.22 and 22.1 ng.g⁻¹) during autumn and spring at Diwaniyah and Hilla River in root samples of both plants respectively. High value of root concentration factor (RCF) had been recorded for HMW (11.7 and 4.46) during the summer at Diwaniyah and Daghara Rivers whereas low value was detected to LMW (0.38 and 0.94) during the summer at Diwaniyah River St.4 and St.5 for both plants respectively. Values of translocation factor were recorded (0.62 and 16.7) of LMW-PAHs for both plants respectively, during spring whereas low factors were recorded (0.03 and 0.1) during summer in both plants. Current study manifested that *P. australis* and *T. domngnesis* were effective in dissipation of PAHs from sediment by the accumulation of HMW-PAHs and dissipation of LMW-PAHs in the roots. Accordingly, these aquatic plants might be useful in evaluating biodegradation of PAHs in freshwater sediment.

Key words: Root concentration factor, Translocation factor, Diwaniyah River, Biodegradation.

عنوان

مجلة العلوم الزراعية العراقية – 47(2): 656-666 / 2016

كفاءة جذور نباتي القصب *Phragmites australis* والبردي *Typha domingensis* في ازالة المركبات الهيدروكربونية الاروماتية متعددة الحلقات من رواسب المياه العذبة

سعد والي علوان

مدرس

كلية الزراعة/جامعة القادسية

abo_hussian@yahoo.com

المستخلص

نفذت الدراسة لقياس تراكيز المركبات الهيدروكربونية الأروماتية متعددة الحلقات (PAHs) في جذور نباتي القصب *Phragmites australis* والبردي *Typha domingensis* وتقييم كفاءتها في معالجة هذه الملوثات من رواسب المياه العذبة في انهار الحلة والديوانية والدغارة. تم جمع عينات الدراسة موسميا خلال المدة (2011-2012). وأظهرت النتائج كفاءة جذور نباتي القصب والبردي في تراكم المركبات ذات الاوزان الجزيئية العالية HMW-PAHs التي كانت تساوي (69.9 و 35.67 نانوغرام.غم⁻¹ وزن جاف) خلال فصلي الشتاء والربيع بينما بلغت (21.8 و 38.7 نانوغرام.غم⁻¹) في عينات الاوراق خلال الصيف والخريف لكلا النباتين على التوالي. وفي مقابل ذلك فان المركبات ذات الاوزان الجزيئية الواطنة-LMW PAHs لم يتم الكشف عنها في بعض المواسم والمحطات وبلغت اعلى هذه القيم (4.22 و 22.1 نانوغرام.غم⁻¹) خلال فصلي الخريف والربيع في نهري الديوانية والحلة في عينات جذور النباتات المدروسة على التوالي. بينت الدراسة ارتفاعاً في قيم معامل تركيز الجذور (Root concentration factor) بين (4.46 و 11.7) للمركبات عالية الوزن الجزيئي HMW-PAH خلال فصل الصيف في نهري الديوانية والدغارة بينما وصلت اقل القيم (0.38 و 0.94) للمركبات الواطنة الوزن الجزيئي خلال فصل الصيف في نهر الديوانية لكل نبات على التوالي. وبلغت قيم معامل النقل (0.62 و 16.7) لمركبات ال LMW - لكل نبات على التوالي خلال فصل الربيع بينما كانت العوامل منخفضة (0.03 و 0.1) خلال فصل الصيف في كل من مركبات ال HMW. تشير الدراسة الحالية الى أن جذور نباتي القصب والبردي لهما القدرة في تراكم وتقييد مركبات HMW-PAHs من الرواسب وتبديد مركبات ال LMA-PAHs لذلك فان هذه النباتات المائية سوف تكون مفيدة في تقييم تحلل وإزالة هذه الملوثات من رواسب المياه العذبة.

الكلمات المفتاحية: معامل التركيز الحياتي للجذور، معامل النقل، نهر الديوانية، التحلل الحيوي

INTRODUCTION

Polycyclic aromatic hydrocarbons (PAHs) are groups of the highly persistent organic pollutants having carcinogenic, mutagenic and teratogenic properties. PAHs have reached into the environment via incomplete combustion of fossil fuels as industrial by-products, which may be dispersed in the air and reach into the water, then accumulate in sediments and soils due to their polarity and lipophilic properties (22). High concentration of PAHs has been found in water, sediment and biota in the Euphrates River and its branches in the middle of Mesopotamia via (34, 2, 4). Many studies referred to the accumulation of the PAHs in plants and their efficiency in the dissipation of them through phytoremediation and rhizosphere microorganism biodegradation. Contribution of plants to PAH dissipation and degradation strongly dependent on the rhizosphere processes (36, 20). Remediation of PAHs in contaminated soil enhances via plants through various processes, such as stimulated on the microbial activity, community structure and functional diversity (42, 24), improvement of soil conditions (3), release of the root exudation (33, 13, 14) and enhancing gas exchange by increasing of the air flow (27,46). The effect of PAHs on wetland plant growth could be species-specific regardless of PAH types and media (52). Decreasing of the dry weight and shoot length of *Phalaris arundinacea* and *Phragmites australis*, and increasing number of young shoots are related to increasing in exposure level of Phenanthrene (23). The accumulative potential of phenanthrene in plant tissues of submersed macrophyte *Vallisneria spiralis* to phenanthrene in freshwater sediment was corresponded to high initial concentrations of phenanthrene in sediments and roots, which was higher than those in the shoots and concentrations of phenanthrene in stolon were lowest among these plant tissues.

Vallisneria spiralis plant could use adaptive mechanisms to toxic contaminants in sediment, and then change the growth patterns to decrease the toxicity on growth, (50). Toyama, *et al.* (45) showed that accelerated removal of pyrene and benzo [a] pyrene in *P. australis* rhizosphere sediments with plants by rhizosphere bacteria. Also, phenolic

compounds in root exudate supported growth as a carbon source and induced benzo [a] pyrene-degrading activity of the mycobacterium. Then, biodegradation of PAHs and the concentration of phenolic compounds increase by stimulation effect in root exudates of *P. australis* exposed to pyrene and benzo(a)pyrene (45). Zhan *et al.* (51) found a relationship between root morphology and compositional factors with plant uptake of Phenanthrene in five crop plant species using a hydroponic system. In soils, the availability of PAHs have been shown to be a primary control for their uptake by plants (29, 18). PAHs tend to strongly partition onto soil solids (particularly the soil organic fractions) due to nonpolar hydrophobic molecules particularly HMW-PAHs. The removal efficiency of PAH differed significantly between LMW-PAHs (phenanthrene 3-rings) and HMW-PAHs (pyrene 4-rings), whereas Pyrene was more persistent in soils than phenanthrene. However, Zhang *et al.* (52) indicated that the growth of wetland plants was significantly influenced by PAHs regardless of the media (water or soil) and PAH types (naphthalene or mixture of phenanthrene and pyrene). Meng *et al.* (33) found that 2-4 ring PAHs were dissipating and losses to be taken up than 5–6-ring PAHs in contaminated soil, probably due to their smaller size increasing their solubility and diffusivity from the soil solution into root cell membranes. Uptake of the organic molecules depending on lipophilicity of the compounds, compositional and morphological feature of roots (37, 51). Therefore, hydrophobic chemicals ($\log K_{ow} > 3.0$) are not easily transported due to the high proportion of lipids located at the cell surface (41). Whereas LMW-PAHs actively transported through plant membranes because of highly water soluble and not sorbed to the roots. So, the size and molecular weight of PAHs may play a role in the ability of a plant to take up the contaminant (38). As well as, the roots and rhizomes of (*P. Australis*) contributes in aeration of the sediment by leakage of oxygen from the roots may create oxidized microzones. (9). The current work aimed to evaluate the root efficiency in PAHs accumulation and to determine root

concentration factor (RCF) and translocation factor (TR) of PAHs in *Phragmites australis* and *Typha domingensis* plants grown in fresh water sediment of Euphrates River branches. In addition, a compositional profile of LMW-PAHs and HMW-PAHs compounds in root of both plants during four seasons were investigated as well.

MATERIALS AND METHODS:

1-Sample Collection and Extraction:

Two species of aquatic plants (*Typha domingensis* Pers.) and (*Phragmites australis*(Cav.) Trin ex Steudel) as the dominant species in Iraqi fresh water were selected in this study, they were classified according to flora of Iraq (44). Plants were collected with their roots from five sites on Diwaniyah and Daghara rivers (Fig.1). Shoots and roots were washed, and aerial dried in the dark and cool cabin. Plants were grounded and homogenized. Five grams of each sample were extracted by ultrasonication for 1 h in 150 ml of acetone and hexane mixture 1:1 (v/v). The solvent was decanted, collected and replenished. Twice solvent fractions were combined and passed through an anhydrous Na₂SO₄ column with elution in 1:1 (v/v) acetone and hexane. The solvents were then evaporated and exchanged to 2 ml hexane, followed by filtration through 2 g silica gel column with elution by 11 ml of 1:1 (v/v) hexane and dichloromethane. Then the organic solvent mixture was evaporated and exchanged to cyclohexane with a final volume of 2 ml (26).

2-PAHs analysis in root:

An external standard method was used in quantifications of 16 PAHs based on five-point calibration curves for individual compounds. Plant samples were analyzed by high performance liquid chromatography Shimadzu (UV/vis. Detector 2500, and supelcosil LC –PAHs column (50 *4. 6 mm, C18, 3µm particles) at the college of Pharmacy lab. / Kufa University. Acetonitrile

and water (both HPLC-grade) as mobile phase in a linear gradient program, (60% acetonitrile (0.3 minute) to 100% (over 2.7 min) with flow rate 2ml/min, Detector (UV, 254) 5 µl of the sample was injected into a capillary column. Peaks in the chromatogram have identified by comparison of the retention time and spectra of standard with those in the sample (1, 16). Standard 16 PAHs were obtained from Aldrich Chemical Co. they are included Naphthalene (Nep) Acenaphthylene (Acpy), Acenaphthene (Acp), Fluoranthene (Flu), Phenanthrene (Phe), Anthracene (Ant), Fluoranthene (Fla), Pyrene (Pyr), Benzo(a)anthracene (B[a]A), Chrysene (Chr), Benzo(b)fluoranthene (B[b]F), Benzo(k)fluoranthene(B[k]F), Benzo(a)pyrene (B[a]P), Dibenzo(a,h) anthracene (D[ah]A), Indeno(1,2,3-c,d)pyrene (InPy), Benzo(g, h,i)perylene (B[ghi]P). Sixteen PAHs have been divided into two different groups, low molecular weight LMW-PAHs have 2-3 rings (Naphthalene to Anthracene) and high molecular weight HMW- PAH have 4-6 rings (Fluoranthene to Indeno(1,2,3-cd) pyrene) (11,10, 35). Root concentration factor (RCF) and translocation factor (TF) were calculated according to the following equations:

$$RCF = \frac{\text{con. of PAH in root (ng.g}^{-1} \text{ dw)}}{\text{con. of PAHs (}\mu\text{g/g) in sediment. (8)}$$

$$TF = SCF / RCF \text{ (33)}$$

Where: SCF is shoot concentration factor.

PAHs concentration data of sediment and shoots of both plants were taken from other work.

3-Statistical Analysis:

Microsoft Excel statistical software and SPSS program were used for analyzing current data in determination of the least significant differences LSD values. Paired T-test were performed at $P < 0.05$ to compare other results.




Fig.1. Map of study region in Al-Qadissyah governorate.

RESULTS AND DISCUSSION:

1-PAHs Concentrations in Plants:

Results showed that high concentrations of HMW-PAHs in *P. australis* roots ranged from 0.2-69.9 ng.g⁻¹ dry weight (DW) during summer at Daghara and Diwanyah River respectively, (table1) whereas LMW-PAHs are not detected during some seasons and stations whereas it had approached 4.22 ng.g⁻¹ during autumn at Diwanyah River. The HMW and LMW in *Typha domingensis* roots were found to be 35.67 and 22.1 ng.g⁻¹ in samples collected in summer from Diwanyah and in the spring at Hilla Rivers respectively, whereas they were not detected

at other stations. Concentrations of LMW ranged from 2.6-44.2 ng.g⁻¹ in *P. australis* shoots in the summer and spring at Hilla and Diwanyah River respectively, whereas HMW ranged 0.27-21.8 ng.g⁻¹ in summer and spring at Daghara and Hilla River respectively. LMW-PAHs level increased from 1.2 to 98.6 in *T. domingensis* shoots whereas HMW concentrations decreased from 0.3-38.7ng.g⁻¹ during spring and autumn at Hilla and Diwanyah River respectively, with significant differences between both groups of compounds (LSD= 15.2 and 3.2) in roots and shoots of *Phragmites australis*, respectively, whereas it was 7.1 and 2.4 in

roots and shoots samples of *T.domingensis*. These results referred to increasing of HMW-PAH concentration in root of both plant on Diwaniyah River, that might be related to either, the high values of total PAHs in sediment of this site which are located under different pollution sources (4) or accelerated biodegradation of LMW through root exudate and associated microorganisms occurred at the rhizosphere (5), or it could be absorbed through plant roots and transported to shoots whereas high molecular weight PAHs can be strongly adsorbed on the root epidermis because of high lipophilicity and resists the biodegradation (49, 26). Also (Günther *et al.* (21), Binet *et al.* (6,7) and Fang *et al.* (17) stated that ryegrass (monocotyledons) had often been selected for the remediation of HMW-PAH in polluted soil because of the increasing in rhizosphere microorganisms. However, PAHs adsorb to plant cell walls, and then they gradually diffuse into subcellular fractions of tissues. Lipid content of intracellular components limited the accumulation of lipophilic compounds, and the diffusion rate is related to the concentration gradient established between cell walls and cell organelles. Unique work showed that Phenanthrene (3 rings) entered the epidermis of the root maize or wheat cells radially and once in the cortex, the apoplasmic flow appeared to dominate the symplasmic movement (48). In addition, many concentrations of PAHs were found in plant roots than in leaves. This plant aided to dissipation of 5- and 6-ring PAHs in soil because of rhizosphere microorganisms activity (6). Generally, according to their molecular weight, plants absorb PAHs either as a gas through the leaves or as a particulate phase by their roots (15, 43). However, emergent plant could absorb semi volatile organic pollutant by two routes, atmospheric deposition, root absorption of PAHs were grown in polluted sediment, then PAHs accumulated in plant roots could transport to the shoots by the transpiration stream (40,19).

However, the relative importance of the stomata and root uptake for contaminant fluxes to leaves is related to the water-solubility of individual PAHs, which increases with decreasing molecular weight. These conclusions might support our results in detecting high concentrations of LMW-PAHs in leaves (Table 1). Li *et al.* (30) was attributed dominance of LMW 2-3 ring, especially naphthalene, in the leaves of *Anatolia. santolin* for absorption of PAHs from air. However, the low concentration of LMW and HMW- PAHs were found during summer because of increasing of photooxidation and volatilization that were related to high temperature in the study region (4). A similar pattern was observed with increasing of HMW-PAH in *Phragmites australis* and *Typha domingensis* roots during the investigation period. This may be related to growing of both species in the same environment and expose to the same pollution sources, also a high accumulation of PAHs in the roots than in the shoot due to poor PAH transport from the roots to aboveground portions of the plant (19). So, Common reed plants might release organic materials by the roots to the surrounding medium (water and bottom sediment), which might encourage the huge amount of bacteria found on the surface of the root system (28). However, *Typha* spp. and *Scirpus lacustris* have been used in horizontal-vertical macrophyte based wetlands to treat phenanthrene.(31). Wei *et al.* (47) indicated that the remediation ability of PAHs attributed to leaching, volatilization, photodegradation, desorption and a biotic factors such plant species, plant metabolism, bioaccumulation, microbial degradation and plant microbial interactions.

Table 1. Seasonal changes in the concentrations of 2, 3, 4, 5 and 6 rings-PAHs compounds (ng.g⁻¹) in roots and shoots of *Phragmites australis* collected from the study stations during (2011-2012).

Stations	PAHs Season	2 rings		3 rings		4 rings		5 rings		6 rings	
		Ro ot	Shoo t	Roo t	Shoo t	Root t	Shoo t	Root t	Shoo t	Roo t	Shoo t
Hilla River	Spring	0.1	35.2	22.1	9.8	22.1	4.12	7.4	1.8	7.17	0.33
	Summer	ND	1.2	ND	2.03	ND	2.03	0.14	1.7	0.1	0.3
	Autumn	0.2	17.3	0.1	14.8	0.1	10.4	1	12.9	1.2	2.56
	Winter	5.5	14.4	0.22	30.05	0.22	20.4	7.3	7.9	8.96	4.4
Daghara River St.2	Spring	0.24	6.32	3.5	52.38	0.2	4.79	0.43	4.8	0	0.86
	Summer	ND	2.3	1.1	3.07	6.48	0.97	8.4	0.97	12	0.81
	Autumn	1	21.3	1.2	71.24	10.3	10.64	2.24	10.67	2.5	3.14
Daghara River St.3	Winter	11.7	26	2.2	42.9	2.6	34.8	20.9	34.8	7.7	6.4
	Spring	0.1	6.3	ND	52.38	4.4	14.6	5.7	4.8	7.5	0.86
	Summer	ND	1	0.11	3.07	0.8	3.8	0.1	0.97	0.1	0.81
	Autumn	13.2	21.3	12.6	71.2	2.4	21.17	0.2	10.6	0.4	3.1
Diwaniya h River St.4	Winter	15.3	26	1.6	42.9	5.9	24.5	7.6	34.86	2.8	6.4
	Spring	1.6	9.1	1.9	10.54	0.46	7.5	1.6	3.75	2.3	1.4
	Summer	0.2	1.8	0.4	4.37	1	3.8	0.1	2.2	1.2	0.97
	Autumn	1.6	9.5	8.2	98.6	2.6	79.2	18.3	29.4	5.22	1.06
Diwaniya h River St.5	Winter	3.7	7.4	10.2	29.93	3.3	10.6	5.43	7.3	17.7	0.74
	Spring	0.6	8.3	1.2	15.3	3.3	9.7	35.67	12.7	22.5	2.2
	Summer	7.6	2.7	2	6.8	4.5	5.5	2.9	1.9	10.6	0.8
	Autumn	2.4	51.1	4.4	87.89	5.8	38.7	4.6	36.6	2.1	12.9
	Winter	2.2	12.9	2.8	51.16	1.5	26	5.62	25.46	4.3	6.7

Table 2. Seasonal changes in the concentrations of 2, 3, 4, 5 and 6 ring-PAHs compounds (ng.g⁻¹) in roots and shoots of *Typha domingensis* collected from the study stations during (2011-2012).

Stations	PAHs Season	2 rings		3 rings		4 rings		5 rings		6 rings	
		Ro ot	Shoo t	Roo t	Shoo t	Root t	Shoo t	Root t	Shoo t	Roo t	Shoo t
Hilla River	Spring	0.1	6.2	ND	23.4	23.45	21.8	6.32	4.1	1.91	1.05
	Summer	0.08	2.6	ND	6.4	12.4	4.8	2.54	1.4	0.98	0.4
	Autumn	0.8	7.8	2.2	24.4	26.6	17.2	4.8	7.2	1.1	2.7
	Winter	1.2	1.8	2.6	3.7	30.4	4.4	6.2	1.7	1.6	0.46
Daghara River St.2	Spring	0.4	11.7	1.8	16.3	22.4	5.8	6.8	2.02	2.2	0.75
	Summer	0.03	2.4	0.1	15.6	0.2	6.07	4.62	1.97	12.9	0.39
	Autumn	0.8	6.01	1.6	17.9	2.2	18.44	10.53	9.8	2.74	1.46
Daghara River St.3	Winter	ND	3.4	1.8	3.4	11.8	5.5	6.81	4.5	38.2	0.57
	Spring	ND	11.7	1.1	25.1	2.2	13.55	3.3	4.11	2.5	1.02
	Summer	0.4	2.3	0.8	4.6	1.2	3.8	28.22	1.2	3.3	0.27
	Autumn	1.8	6.1	ND	9.7	1.4	8.01	9.11	14.46	7.3	0.79
Diwaniya h River St.4	Winter	1.8	3.4	0.8	7.3	1.8	4.9	11.4	3.6	3.2	1.8
	Spring	0.08	6.5	0.2	2.2	12.3	4.86	3.7	3.13	10.3	2.31
	Summer	0.2	1.13	3.5	0.37	68.1	1.6	8.4	0.36	2.1	0.54
	Autumn	0.4	5	0.84	1.66	42.5	10.05	20.7	8.05	9	3.11
Diwaniya h River St.5	Winter	1.6	4	ND	1.3	69.9	3.79	17.2	4.9	15.2	2.9
	Spring	1.4	19.2	1.6	44.2	6.6	6.56	4.4	13.15	2.6	3.06
	Summer	1.1	6.4	0.9	7.66	1.3	7.9	2.2	3.6	2.1	0.36
	Autumn	1.2	7.2	4.22	9.5	42.7	6.7	10.7	7.5	6	0.97
	Winter	2.4	13	1.2	12.22	37.9	5.6	34.8	3.4	11.8	1.2

2-Root Concentration Factor and

Translocation Factors:

Data of the current study indicated that there was a significant increment of root concentration factor (RCF) of HMW-PAHs compared to LMW-PAHs in *P. australis* (Fig.2) and *T. dominance* roots at five stations in the study region during four seasons (Fig.3). The high value of RCF was recorded for HMW to be (11.7 and 4.46) during the summer at Diwaniyah (St.4) and Daghara Rivers (St.2), whereas a low value of LMW to be (0.38 and 0.94) during the summer at Diwaniyah River St.4 and St. 5 for both plants respectively. Statistically, T-test indicated a significant increment in RCF values in *P. australis* and *T. domingensis* roots for LMW-PAHs and HMW-PAHs accumulation ($P < 0.05$). These results manifested the different capability in dissipation efficiency of both groups in the roots of both plants and this may be related to the chemical solubility of these chemicals in water and in lipophilicity of PAHs within the component of root cell which is known to be crucial in the transport within the plant shoots. Similar data were reported by Meng *et al.*, (33) in restricted uptake of 5-6 ring of the plant compared to 2-4 ring of PAHs while the uptake of pyrene was several times greater than of Phenanthrene (3 ring). Subsequently, increasing of RCF with increasing of contact time between root and solution and lipophilicity also the accumulation of these PAHs correlated with their concentrations in soil and plant composition (18, 25). Results of translocation factor (TF) showed that high values of LMW-PAHs were detected for both plants, especially in spring, which was ranged (0.4-48) in *P. australis* at St.4 and St.1 respectively (Fig.4). In *T. domngensis*, translocation factor ranged 0.98-43) in

summer at St.5 and St.1 respectively (Fig.5). However, TF values decreased during the winter for all stations except St.5 that may be due to increased pollution level of PAHs in this site (4). Significant differences were detected ($P < 0.05$) of TF between HMW and LMW PAHs from roots to leaves of both plants respectively. This variation might be associated with different molecular weight and solubility of PAHs or related to plant species. In addition, low values of TF with HMW were found for all stations and seasons, whereas high values of them (0.62 and 16.7) were observed on both plants respectively in the spring and low factors of (0.03 and 0.1) noticed in the summer for both plants this can be due to increasing of biodegradation in rhizosphere zone. Also, seasonal variation of organic pollutants probably due to the increasing number of bacteria grown on the root system and bottom sediments. (28). Consequently, the observed difference between *P. australis* and *T. domingensis* in concentrations of PAHs can be linked to plant species characteristics, including rhizospheric properties (root structure, root exudates, presence of rhizospheric microorganisms, rhizobia symbionts). In addition, the transpiration rate and the lipid content of root cell fractions are the main drivers of the subcellular partition of PAHs in roots (25). This work outlined the importance of aquatic plant in promoting the accelerated removal of PAHs in freshwater sediments either by degradation of LMW-PAHs with rhizosphere microorganism or restricted of HMW-PAHs in root tissues. *P. australis* and *T. domngensis* indicated high efficiency in dissipation of PAHs from sediment. These plants have a high ability to tolerate high levels of organic pollutants. So, it can be used as phytoremedidant.




Fig.2. Root Concentration Factor (RCF) of HMW-PAHs and LMW-PAHs in *P. australis* roots in study region during the four seasons (2011-2012). ($P < 0.05$).




Fig.3. Root Concentration Factor (RCF) of HMW-PAHs and LMW-PAHs in *T. domingnesis* roots in study region during the four seasons (2011-2012). ($P < 0.05$).




Fig.4. Translocation Factor (TF) of HMW and LMW in *P. australis* roots in study region during the four seasons (2011-2012).




Fig.5. Translocation Factor (TF) of HMW and LMW in *T. domingnesis* roots in study region during the four seasons (2011-2012).

REFERENCES

- 1- A.P.H.A. (American Public Health Association). 2003. Standard Methods for examination of water and wastewater. 20th Ed. Washington, DC, USA. (3): 6440 A.
- 2- AL-Azaway, A .S. 2012. Biogeochemical distribution of some Polycyclic Aromatic Hydrocarbons(PAHs) in Hilla River-Iraq. Ph.D dissertation. University of Babylon.pp:265.
- 3- Anderson, T.A., E.A Guthrie and B.T Walton. 1993. Bioremediation in the rhizosphere. Environ Sci Technol 27:2630–2637.
- 4- Alwan, S.W. 2013. Ecological Study of Polycyclic Aromatic Hydrocarbons in Diwanayah and Daghara Rivers and their accumulation in some aquatic plants/Iraq. Ph.D dissertation. University of Babylon.pp:195.
- 5- Aprill, W. and R. C. Sims.1990 Evaluation of the use of prairie grasses of simulating polycyclic aromatic hydrocarbon treatment in soil. Chemosphere. 20 (1-2): 253-265.
- 6- Binet, P., J.M Portal and C. Leyval. 2000a. Dissipation of 3-6-ring polycyclic aromatic hydrocarbons in the rhizosphere of ryegrass. Soil Biol. Biochem 32, 2011–2017.
- 7- Binet, P., J.M Portal, and C. Leyval. 2000b. Fate of polycyclic aromatic hydrocarbons (PAH) in the rhizosphere and mycorrhizosphere of ryegrass. Plant Soil 227, 207-213.
- 8- Briggs, G.G., Bromilow, R.H. and Evans, A. A.1982. Relationships between lipophilicity and root uptake and translocation of non-ionized chemicals by barley. Pestic Sci, 13:495-504.
- 9- Brix,H. and H. Schierup. 2011. The use of aquatic macrophytes in water-pollution control. Royal Swedish Academy of Sciences. 18(2): 100-107.
- 10- Brown, J. 2002. Partitioning of chemical urban stormwater. Dissertation, University of Otago, Dunedin, New Zealand.
- 11- Brown, J., & B. Peake. 2006. Sources heavy metals and polycyclic aromatic hydrocarbons in urban stormwater runoff. Science of the Total Environment, 359, 145–155.
- 12- Chen B, J.L. Schnoor .2009. Role of suberan, and hemicellulose in phenanthrene sorption by root tissue fractions of switch grass (*Panicum virginatum*) seedlings. Environ Sci Technol. 43:4130–4136.
- 13- Child R ,C.D. Miller, Y. Liang, G. Narasimham, J. Chatterton, R.C. Sims, D Britt and A.J. Anderson. 2007a. Polycyclic aromatic hydrocarbon degrading Mycobacterium isolates: their association with plant roots. Appl Microbiol Biotechnol 75:655–663.
- 14- Child R, C.D. Miller, Y. Liang, R.C. Sims and A.J. Anderson. 2007b. Pyrene mineralization by Mycobacterium sp strain

- KMS in a Barley rhizosphere. *J Environ Qual* 36(5):1260–1265.
- 15- Edwards, NT. 1988. Assimilation and metabolism of polycyclic aromatic hydrocarbons by vegetation – an approach to this controversial issue and suggestions for future research, pp. 211–229 Cook & Dennis Polycyclic Aromatic Hydrocarbons: A Decade of Progress. 10th Int. Symp, Battelle Press, Columbus, OH Ed. Book.
- 16- Environmental Protection Agency (EPA). 2012. Methods for organic chemical analysis of municipal and industrial wastewater. Appendix A to part 136.(electronic version).
- 17- Fang, C., M. Radosevich and J.J. Fuhrmann. 2001. Atrazine and phenanthrene degradation in grass rhizosphere soil. *Soil Biol Biochem* 33, 671–678.
- 18- Gao, Y.Z., L. Z. Zhu, C.J. Hu and B.L.Chen. 2004. Effects of Tween 80 on plant uptake of phenanthrene and pyrene from water. *Acta Sci Circumstantiae* 24:713–718.
- 19- Gao, Y.Z. and L.Z. Zhu. 2004. Plant uptake, accumulation and translocation of phenanthrene and pyrene in soils. *Chemosphere* 55:1169-1178.
- 20- Gao, Y., S.C. Wu, X.Z. Yu and M. Wong, 2010. Dissipation gradients of phenanthrene and pyrene in the rice rhizosphere. *Environmental Pollution* 158: 2596-2603.
- 21- Gunther, T., U. Dornberger and W. Fritsche 1996. Effect of ryegrass on biodegradation of hydrocarbons in soil. *Chemosphere* 33, 203–215.
- 22- Haritash, A.K. and C.P. Kaushik. 2009. Biodegradation aspects of polycyclic aromatic hydrocarbons (PAHs): a review. *J Hazard. Mater.* 169:1–15.
- 23- Hubner, T.S. , H. Tischer, F. Tanneberg and. Kuschk .2000. Influence of phenol and phenanthrene on the growth of *Phalaris arundinacea* and *Phragmites australis*, *Int. J. Phytorem.* 2: 331–342.
- 24- Jennifer, L.K., N.K John, L. Hung and T.T. Jack. 2005. The effects of perennial ryegrass and alfalfa on microbial abundance and diversity in petroleum-contaminated soil. *Environ Pollut.*133:455–465.
- 25- Kang, F, D. Chen, Y. Gao and Y. Zhang. 2010. Distribution of polycyclic aromatic hydrocarbons in subcellular root tissues of ryegrass (*Lolium multiflorum* Lam.).*BMC plant biology*, 10:210.
- 26- Kipopoulou, A.M., E. Manoli and C. Samara. 1999. Bioconcentration of PAHs in vegetables grown in an industrial area. *Environmental Pollution* 106:369–380.
- 27- Khan, A.A., R.F. Wang, W.W.Cao., D. R. Doerge, D. Wennerstrom and C.E. Cerniglia. 2001. Molecular cloning, nucleotide sequence, and expression of genes encoding a polycyclic aromatic ring dioxygenase from *Mycobacterium* sp strain PYR-1. *Appl Microbiol Biotechnol* 67:3577–3585.
- 28- Lalke-Porczyk, E. and W. Donderski. 2005. The role of bacteria growing on the root system of the common reed (*Phragmites australis* [Cav.] Trin.ex Steudel) in the metabolism of organic compounds. *Polish Journal of Environmental Studies* 14(1): 57-64.
- 29- Li, Y., A.Yediler, Z. Ou, I. Conrad and A. Kettrup. 2001. Effects of a nonionic surfactant (Tween80) on the mineralization, metabolism and uptake of phenanthrene in wheat–solution–lava microcosm. *Chemosphere* 45:67–75.
- 30- Li, J., X. Shanga., Z. Zhaoa, R.L.Tanguaya, Q. Tonga and C. Huang. 2010. Polycyclic aromatic hydrocarbons in water, sediment, soil, and plants of the Aojiang River waterway in Wenzhou, China. *Journal of Hazardous Materials*, 173: 75–81.
- 31- Machate, H.T., H. Noll, Behrens and A. Kettrup.1997. Degradation of phenanthrene and hydraulic characteristics in a constructed wetland, *Water Res.* 31 (3):554–560.
- 32- Meagher, R. B. 2000. Phytoremediation of toxic elemental and organic pollutants, *Curr. Opin. Plant Biol.* 3 : 153–162.
- 33- Meng, L., M. Qiaom, H. Peter and H. Arp. 2011. Phytoremediation efficiency of a PAH-contaminated industrial soil using ryegrass, white clover, and celery as mono- and mixed cultures. *J Soil sediments.* 11:482–490.
- 34- Mohamed, A. B. 2007. Qualitative and Quantitative Studies of some Polycyclic Aromatic Hydrocarbons (PAHs) and Limnology of Euphrates River from Al-Hindiya Barrage to Al-Kifil city – Iraq. PhD dissertation College of science. University of Babylon. pp:240.

- 35- Neff, J.M. 1979. Polycyclic Aromatic Hydrocarbons in the aquatic environment, sources, fates and biological effects. Applied Science, London, UK.
- 36- Pilon-Smits, E. 2005. Phytoremediation. *Ann Rev Plant Biol* 56:15–39.
- 37- Prasad, M.N. 2005. Aquatic Plants for Phytotechnology.. In: Prasad MNV (ed) Marcel Dekker Inc. New York, 259p.
- 38- Ryan, K. and M.K. Miya. 2001. Enhanced phenanthrene biodegradation in soil by slender oat root exudates and root debris. *J Environ Qual*.30:1911–1918.
- 39- Schnoor, J. L., L.A. Licht, S.C. McCutcheon, N. L. Wolfe and L.H. Carreira, 1995. Phytoremediation of organic and nutrient contaminants. *Environmental Science and Technology*. 29 (7): 318-323.
- 40- Sicbaldi, F., G.A. Sacchi, M. Trevisan and A.A. Del Re. 1997. Root uptake and xylem translocation of pesticides from different chemical classes. *Pesticide Science* 50 (2), 111-119.
- 41- Siciliano, S. D. and J.J. Germida. 1998b. Mechanisms of phytoremediation: biochemical and ecological interactions between plants and bacteria. *Environmental Review*. 6: 65-79.
- 42- Siciliano, S., J.J. Germida, K. Banks and C.W. Greer. 2003. Changes in microbial community composition and function during a polyaromatic hydrocarbon phytoremediation field trial. *Appl Environ Microbiol* 69:483–489.
- 43- Simonich, S.L. and R.A. Hites. 1995. Organic pollutant accumulation in vegetation. *Environ Sci Technol* 29, 2905–2914.
- 44- Towensend, C.C. and E. Guest. 1985. Flora of Iraq. Ministry of Agriculture and Agrarian Reform, Vol 9,10. Republic of Iraq.
- 45- Toyama, T.T., T. Furukawa, N. Maeda, D. Inoue, K. Sei, K. Mori, S. Kikuchi and M. Ike. 2011. Accelerated biodegradation of pyrene and benzo[a]pyrene in the *Phragmites australis* rhizosphere by bacteria- root exudate interactions. *Water research* 45: 1629-1638.
- 46- Tsao, D.T. 2003. Overview of phytotechnologies. *Adv Biochem EngBiotech* 78:4–50.
- 47- Wei, S. & P. Shengwang. 2010. Phytoremediation of soils contaminated by phenanthrene and pyrene with multiple plant species. *J. Soils. Sediments*. 10:886–894.
- 48- Wild, E., J. Dent, G.O Thomas and K.C. Jones. 2005. Direct observation of organic contaminant uptake, storage, and metabolism within plant roots. *Environ Sci Technol*. 39:3695–3702.
- 49- Wild, S.R. and K.C. Jones. 1992. Organic chemicals entering agricultural soils in sewage sludges: screening of their potential to transfer to crop plants and livestock. *Sci Total Environ* 119, 85–119.
- 50- Yana, Z.S., H. Guob, T. Songa, Y. Hua and H. Jianga. 2011. Tolerance and remedial function of rooted submersed macrophyte *Vallisneria spiralis* to phenanthrene in freshwater sediments. *Ecological Engineering* 37: 123–127.
- 51- Zhan, Z., X. Liang, G. Xu and L. Zhou. 2013. Influence of plant root morphology and tissue composition on phenanthrene uptake: Stepwise multiple linear regression analysis. *Environmental Pollution* 179. 294-300.
- 52- Zhang, Z., Rengel,Z. and K. Money. 2010. Polynuclear aromatic hydrocarbons (PAHs) differentially influences growth of various emergent wetland species, *Journal of Hazardous Material*. 182:689–695.