

EFFECT OF DIFFERENT FEEDING LEVELS ON PHYSICO-CHEMICAL TRAITS FOR MAIN CARCASS CUTS OF PEKIN DUCKS IN KURDISTAN REGION OF IRAQ

Azad Sh. S. Al-Dabbagh^{1*}Sardar. Y. T. Al- Sardary²

Lecturer

Prof.

Dept. Anim. Reso. – Coll. Agric. Engine. Sci. - Salahaddin University - Erbil, Iraq

E-mail: azad.salih2@su.edu.krd

ABSTRACT

The objective of this study was to investigate the effect of different feeding levels (0%, 25%, and 50%) of alfalfa pellet ration (APR) (47% alfalfa) on the main carcass cuts Physico-chemical traits of Pekin ducks (*Anas platyrhynchos*). A total of 270 one-day-old ducklings were used. The birds at seven-day-old were assigned to 9 indoor pens of (2.2×1m) with outdoor and swimming pool (9.8×1m). Each pen represented an indoor replicate with 30 ducklings. The feeding program for Pekin Duck group treatments with (APR) (47% alfalfa) was used as a replacement to the basal diet at different percentages (0%, 25%, and 50%) for control (T0), first (T1) and second (T2) groups, respectively. At the end of the rearing period at 49 days, 6 birds (3male+3female) were randomly selected from each treatment, slaughtered and each breast and thigh muscle were excised for meat quality analysis. The results showed that treatments and sex did not significantly influence meat Cholesterol content, and most results for each of cooking loss, color, pH, chemical composition, and amino acid profile of the meat. In contrast, the treatments and sex had a significant effect on most for each fat-soluble vitamins, water-soluble vitamins, and fatty acid profiles. In general, feeding ducks with a 25% of APR improved the color of breast, and fat-soluble vitamins with Cholesterol in both breast and thigh muscle while feeding ducks with a 50% APR improved most Physico-chemical traits.

Keywords: Vitamins, breast, thigh, ration, fatty acid, amino acid, birds

*Part of Ph.D. Dissertation of the 1st author.

الدباغ والسرداري

مجلة العلوم الزراعية العراقية- 2025 :56(1):375-390

تأثير مستويات مختلفة من التغذية على الصفات الفيزيوكيميائية للأجزاء الرئيسية في الذبيحة لبط البكين في إقليم كردستان العراق

سردار ياسين طه السرداري²نازاد شمس الدين صالح الدباغ^{1*}

أستاذ

مدرس

قسم الثروة الحيوانية / كلية علوم الهندسة الزراعية / جامعة صلاح الدين – أربيل، العراق

المستخلص

استهدفت هذه الدراسة إلى معرفة تأثير مستويات التغذية المختلفة لعلف أقراص الجت على الصفات الفيزيوكيميائية للأجزاء الرئيسية في الذبيحة لبط البكين. تم إحضار 270 فرخة عمرها يوم واحد من مفسس مزرعة نوروز – أربيل. تم تخصيص 9 حظائر داخلية مغلقة بمساحة (2.2×1م) لأفراخ عمر 7 أيام ومع مساحات خارج الحضيرة وحوض السباحة (9.8×1م)، وكان كل حضيرة يمثل تكراراً داخلياً به 30 فرخة بط. وتم اتباع برنامج التغذية لمعاملات مجموعة البت البكين باستخدام علف أقراص الجت (47% الجت) كإحلال من الغذاء الأساسي وبنسبة مئوية مختلفة (0%، 25% و50%) لثلاث مجموعات على التوالي. في نهاية فترة التربية، تم اختيار 6 طيور (3 ذكور+3 أنثى) بشكل عشوائي من كل معاملة وذبحها، وتم استئصال كل من عضلات الصدر والفخذ لتحليل جودة اللحوم. أظهرت النتائج أن المعاملات والجنس لم تؤثر معنوياً على محتوى الكوليسترول ومعظم النتائج الخاصة بفقدان الماء أثناء الطبخ، اللون، قيم للأس الهيدروجيني، التحليل الكيميائي والأحماض الأمينية، وفي المقابل كان للمعاملات والجنس تأثير معنوي على محتوى معظم الفيتامينات الذائبة في الماء والدهون والأحماض الدهنية. بشكل عام، تغذية البت بنسبة 25% من علف أقراص الجت أدت إلى تحسين لون الصدر، والفيتامينات الذائبة في الدهون مع الكوليسترول في كلا الصدر و الفخذ، في حين أن تغذية البت بنسبة 50% من علف أقراص الجت أدت إلى تحسين معظم الصفات الفيزيوكيميائية.

الكلمات المفتاحية: الفيتامينات، الصدر، الفخذ، العلف، الأحماض الدهنية، الأحماض الأمينية، الطيور

* 1 جزء من بحث أطروحة دكتوراه للباحث الأول

INTRODUCTION

The Pekin duck was created in the second half of the nineteenth century. Commercial Pekin ducks are most often reared for meat and have snow-white plumage (35). These ducks are characterized by a fast growth rate, high slaughter weight, good feed conversion, high viability, relatively low nutrient requirements compared to broiler chickens, great tolerance to harsh raising conditions, and considerable to disease immunity (17, 25). The number one duck meat producer was China, which produced about two-thirds of the total world production (10). Pekin-type broiler ducks are raised intensively from 6 to 7 weeks, and under a backyard system from 9 to 10 weeks. Ducks are a great choice for backyard and organic farming for the reasons mentioned above as well as the excellent conversion of farm feeds (forages, feed roots, dried feeds, silages) and by-product feeds (5). The Pekin-type duck carcass is a favorite among poultry farmers and is distinguished by having more red muscle fibers in the breast muscle than chicken meat. In general, duck meat is thought to be flavorful, high in polyunsaturated fatty acids, and amino acids, and relatively little fat. With respect to the overall content of polyunsaturated fatty acids, duck meat has a high content of linoleic and linolenic fatty acids (27). Furthermore, Alfalfa (*Medicago sativa* L.) in ducks feed, is primarily grown for and used in animal feed, as it is a good source of easily assimilated protein and high in minerals and vitamin content. Based on this potential, alfalfa pellet ration can be utilized as a food replacement to reduce cholesterol and enhance the quality of duck meat. Duck meat has also a favorable profile of essential amino acids (including relatively high amounts of lysine) and the highest content of unsaturated fatty acids among all poultry species. It is a source of fat-soluble vitamins (mainly A and E) and water-soluble vitamins (B1, B2, and PP) (16). The aim of this study was to investigate the effect of different alfalfa pellet ration levels on Physico-chemical traits for the main carcass cuts breast and thigh of Pekin ducks in Kurdistan region of Iraq.

MATERIAL AND METHODS

Management and Design of Experiment:

This study was conducted at the private field

Nowruz farm - Qushtapa - Erbil during the period from 9 Sept. 2020 until 11 November 2020 using different feeding levels for Pekin Duck (*Anas platyrhynchos*). A total of 270 one-day-old ducklings were brought from Nowruz farms hatchery – Erbil. The birds at seven-days-old were assigned to 9 indoor pens (2.2×1m) with outdoor and separated swimming pools (9.8×1m); each box represented an indoor replicate with 30 ducklings. The feeding program for Pekin Duck group treatments about (starter, growing, and finisher), and alfalfa pellets ration (47% alfalfa) were used at different levels (0%, 25%, and 50%) for three groups respectively, feed and water were provided ad libitum. Basal diet ingredients, Alfalfa pellet ration ingredients, and calculated chemical composition of the mixed them are shown in Table (1, 2, 3), respectively. The duckling was in a clean well-ventilated hall and belonged to a regular healthy program applied on the farm. All ducklings were vaccinated (Table 4). In the first week, the ducklings were fed collectively ad libitum on the starter full requirements of the Basal Diet for broiler feed reared indoor house according to NRC (26) only and freely in a hall where the number of lighting hours was about 22 hours, and with an area of 27 birds / m². Vitamin C was given about 1 g/ 1 liter of water after vaccination of the ducklings with Newcastle and influenza on three consecutive days. The multi-vitamin was also given by drinking water 1 ml / 1 liter of water three times a day for three consecutive days during the first week of the experiment. To improve the amount of feed intake, the alfalfa pellet ration was crushed and presented during the first week of the experiment. Also, 6 ml / 1 liter water di-calcium phosphate was given in liquid form, from 2 to 4 weeks of age, for three consecutive days each week. Birds were randomly divided into three treatment groups of three replicates / treatment of 30 birds / replicate at one week's age, and then different dyeing color was used to label each treatment. The treatments were as follows; control treatment group: as control (T0) fed full requirements of basal diet according to NRC (26) using broiler feed, first treatment (T1): birds fed %25 alfalfa (*Medicago sativa*) pellet ration + 75% of full requirements of basal diet,

second treatment (T2): birds fed %50 alfalfa pellet ration + 50% of full requirements of basal diet. From 15 days, birds are enabled to reach the swimming pool (10m×1m×30cm) through separate holes from indoor to outdoor. The measures of the area in indoor and outdoor with swimming pool 12 × 1m (1 m²/4 bird) for each pen and separated by fences.

The outdoor area was designed to provide birds with natural behavior. All birds were supplied with clean water ad libitum as well as a clean water pool. At the end of the rearing period at 49 days, 6 birds (3male+3female) were randomly selected from each treatment, slaughtered and each breast and thigh muscle were excised for meat quality analysis.

Table 1. Basal Diet ingredients (%).

Ingredients	Starter	Grower	Finisher
	1-14 days	15-35 days	36-49 days
Wheat	19.5	23	18.6
Wheat bran	1	2.5	4
Wheat flour	10	10	15.5
Corn	23	24	30.5
Soybean	39.85	34.3	24.7
Oil	1.85	1.5	2
Limestone	1.9	1.8	1.8
Premix	2.5	2.5	2.5
Optifeed*	0.005	0.005	0.005
Oleobiotec**	0.005	0.005	0.005
VeO***	0.005	0.005	0.005
Lysin	0.02	0.03	0.04
Methionine	0.05	0.05	0.05
Avimatrix****	0.05	0.05	0.05
Herb – All (COCC-X) *****	0.05	0.05	0.05
Antitoxin	0.075	0.075	0.075
Chemical composition (%)			
Protein	23	22	18.5
Metabolic Energy(Kcal/Kg)	2850-2900	2900-3000	3000-3100
Moisture	11	10.9	10.5
Fat	3.4	3.8	3.5
Fiber	3.6	3.5	4.0
Ash	5.1	5.0	4.8

* It is a flavoring substance for stimulation of the appetite.

** It is a mixture of essential oils and spices to support the functioning of the microbiota and increase digestive secretions and gastric motility.

***It is a blend of natural and purified fractions from plant extracts that regulates appetite and reduce stress.

**** It also contributes to improving feed conversion and optimal production efficiency.

***** It is a complementary feed to improve and strengthen gut health for all animal species that stimulate appetite

Table 2. Alfalfa pellet ration ingredients (%).

Ingredients	%
Alfalfa	47
Soybean	8.6
Wheat bran	2.5
Wheat flour	33.6
Oil	2
Sodium carbonate	0.3
Antitoxin	0.1
Dicalcium phosphate	0.5
Limestone	2
Salt	1
Premix (vitamins, Lysin, Methionine, and minerals)	1
Dates juice	1.5
Chemical composition (%)	
Protein	15.9
Metabolic Energy (Kcal/Kg)	2400 - 2600
Moisture	12.0
Fat	1.3
Fiber	8.8
Ash	8.1

Table 3. Calculated chemical composition (%) of the mixed Basal diet and Alfalfa pellet rations.

Chemical composition (%)	Starter 7-14 days			Grower 15-35 days			Finisher 36-49 days		
	T0	T1	T2	T0	T1	T2	T0	T1	T2
	Control BD* (100%)	APR** (25%) + BD (75%)	APR (50%) + BD (50%)	Control BD (100%)	APR (25%) + BD (75%)	APR (50%) + BD (50%)	Control BD (100%)	APR (25%) + BD (75%)	APR (50%) + BD (50%)
Protein	23	21.225	19.45	22	20.475	18.95	18.5	17.85	17.2
Metabolic Energy (Kcal/Kg)	2850- 2900	2737.5- 2825	2625- 2750	2900- 3000	2775- 2900	2650- 2800	3000- 3100	2850- 2975	2700- 2850
Moisture	11	11.25	11.5	10.9	11.175	11.45	10.5	10.875	11.25
Fat	3.4	2.875	2.35	3.8	3.175	2.55	3.5	2.95	2.4
Fiber	3.6	4.9	6.2	3.5	4.825	6.15	4.0	5.2	6.4
Ash	5.1	5.85	6.6	5.0	5.775	6.55	4.8	5.625	6.45

* BD: Basal Diet ** APR: Alfalfa Pellet Ration T0: Control

T1: First treatment T2: Second treatment

Health control of Pekin duck

The duckling was in a clean well-ventilated hall and belonged to a regular health program. All ducklings were vaccinated based on production purposes.

Table 4. Vaccination program for Pekin duckling

Age Day	Vaccine	Route
3	IB Ma5+ ND Clone 30	Spray
7	InfluenzaH9N2+ ND	Injection

2.2. Evaluation of meat Physico-chemical traits

2.2.1. Cooking loss

The muscle samples from the breast and thigh were transferred from the -18°C freezer into a 4°C cooler overnight to thaw. The defrosted samples were individually weighed and documented as initial weight (W1). The samples were then placed in plastic bags and cooked in a water bath for 20 minutes at 80°C. The cooked samples were removed from the plastic bags and cooled in an ice slurry for 20 minutes. The samples were weighed once again and then written as W2 (cooked weight). The following equation was used to determine

the cooking loss based on the weight differences between raw and cooked meat:

$$\text{Cooking loss (\%)} = ((W1 - W2) \div W1) \times 100$$

According to the method recommended by Abdullah et al. (1).

2.2.2. Color measurement

Breast and thigh samples were taken out of the -18°C freezer and allowed to defrost overnight at 4°C . Prior to color measurement, they were taken out of the packaging and given 20 minutes to bloom in the air. A Color Flex Spectrophotometer (Shenzhen 3nh Technology Co., Ltd, China) was used to measure the color of the meat. Prior to use, the device was initially calibrated against black-and-white reference tiles. On each sample, L^* (lightness), a^* (redness), and b^* (yellowness) were measured in triplicate (1).

2.2.3. pH muscle determination

A digital pH meter was used to indirectly measure the pH of the breast and thigh muscles under three post-mortem conditions: pH_{60} (one-hour post-slaughter), pH_{24} (24-hour post-mortem), and pH_f (post-freezing) (Electronic Co. Ltd, China). Before each use, the pH meter's calibration was done at pH 4.0 and pH 6.8. Each crushed muscle sample weighed around 0.5 g, which was then homogenized for 20 seconds in 10 ml of ice-cold double distilled water. The pH of homogenates was determined using the pH meter's electrode. Each sample was measured three times, and for each treatment, the average pH value was calculated., and also according to the technique suggested by Abdullah et al. (1).

2.2.4. Proximate composition

Following AOAC (3) guidelines, the breast and thigh meat's composition was identified. Moisture was calculated by drying 50 g of meat at $100\text{-}105^{\circ}\text{C}$ until it attained a constant weight. The Kjeldahl method was used to calculate crude protein. The crude protein was gained as $6.25 \times N\%$. The Soxhlet extraction technique and petroleum ether were used to calculate the meat's fat content. By burning the sample at 550°C for 3 hours in a muffle furnace, the amount of Ash in the meat was determined.

2.2.5. Vitamins

Fat soluble vitamins A, D, E, K were determined by using the liquid

chromatography technique (HPLC) model (SYKAMN – Germany) in the laboratory of the Ministry of Science and Technology – Baghdad - Iraq, the column separation was (C18 – ODS (25 cm * 4.6 mm) and the mobile phase was methanol–water (98:2, v/v). The mobile phase flow rate was 1.0 mL/min, and the injection volume was 50 μL . The column oven temperature was 35°C . Detection with a UV–vis detector was carried out at 265 nm According to the procedure mentioned by Xue et al., (39). Water soluble vitamins (B2, B3, B5, B6, B12, C) were determined by using HPLC model SYKAMN (Germany) It was used to analyses add detection of thiamethoxam. The mobile phase was an isocratic acetonitrile: D.W: formic acid (50: 47: 3) at flow rate at 1.0 mL/min, column was C18 – ODS (25 cm * 4.6 mm) and the detector UV- 215, 210 nm according to (33).

2.2.6. Fatty acid profile and cholesterol

Fatty acid was determined by using the Gas chromatography (GC-2010) model (Shimadzu – Japan) in the laboratory of the Ministry of Science and Technology – Baghdad - Iraq, the column separation was SE- 30) (30m * 0.25 mm), and it used Flame Ionization Detector (FID). The gas flow rate was 100 Kpa (Kilo pascal). The injection temperature was 280°C and indicator temperature was 310°C . The separator column temperature was $120 - 290^{\circ}\text{C}$ ($10^{\circ}\text{C} / \text{MIN}$), According to (40). Cholesterol in breast and thigh samples was extracted and determined by HPLC and using equipment as described in Maiorano et al. (23).

2.2.7. Amino acid profile

The amino acid was determined by using the amino acid analyzer-Korea in the laboratory of the Ministry of Science and Technology – Baghdad - Iraq, the column separation was ZORBAX Eclipse-AAA; $3.5\mu\text{m}$; $L \times i.d.=150 \times 4.6 \text{ mm}$ to separate amino acids. The mobile phase was acetonitrile: methanol: formic acid (60: 20: 20) rates with flow rate 1 ml / 1 min. The gas flow rate was 100 Kpa (Kilo pascal). The injection temperature was 280°C and indicator temperature was 310°C . The separator column temperature was $120 - 290^{\circ}\text{C}$ ($10^{\circ}\text{C} / \text{MIN}$), According to Dahl-Lassen et al. (7).

2.3. Statistical Analysis: The experiment was designed as Complete Randomized Design (CRD). Data were analyzed using the program of Statistical Analysis System (32), The effect of sex (male or female) was added to the module of all traits. To diagnosing the significant differences between treatments, the proceeding of Duncan's multiple range tests at level of $P \leq 0.05$ was detected (9).

RESULTS AND DISCUSSION

3.1. Cooking loss

The effect of the replacement of alfalfa pellet to the basal diet on cooking loss in Pekin duck's breast and thigh meat is shown in Table 5. Irrespective of the treatment, no significant differences ($P > 0.05$) on the cooking loss was observed in breast muscles among birds fed the control and those fed basal diets with replacement alfalfa pellet after freezing storage, and the first treatment (T1) gave higher cooking loss (30.15%) than in the second treatment (T2) (28.57%) and control (T0) (26.39%) for breast respectively, while control (T0) (28.52%) and (T1) (26.64%) was higher significantly ($P \leq 0.05$) than (T2) (20.31%) for thigh respectively. Regarding sex effect, the cooking loss was lower ($P > 0.05$) in the breast of males than in females, and high in the thigh of females than in males (Table 5). Addini et al., (2) reported similar findings where birds fed alfalfa supplementation of 5% and 10% had higher cooking loss% than the control of hybrid duck meat. However, He et al., (15) referred to similar findings too in their study on broilers where fed alfalfa meal supplementation of 25g/kg, 50g/kg, and 75g/kg feed had no significant effect on cooking loss% in comparison with the control in the breast but different findings in the thigh which showed no significant differences on cooking loss%. In contrast, Kokoszynski et al. (18), showed in their study on Pekin duck and commercial crossbreds that sex effect on breast and legs, females recorded lower than males of Pekin ducks in the breast but similar results in leg that males had lower cooking loss than females in commercial hybrid Pekin ducks. However, other studies referred that those of males and females had no significant differences ($P > 0.05$) in breast and leg (21, 20).

Table 5. Effect of treatments and sex on cooking loss (%) of breast and thigh for Pekin ducks

Factors	Breast	Thigh
Over all mean	28.37	25.16
Treatments	NS	*
T0	26.39±1.25	28.52±0.31a
T1	30.15±1.54	26.64±1.09a
T2	28.57±1.16	20.31±0.83b
Sex	NS	NS
M	27.83±1.38	25.69±2.28
F	28.91±1.39	24.62±2.74

Means followed by the different letters in the same column are significantly different ($P \leq 0.05$). *: $P \leq 0.05$ NS: non-significant

3.2. Color values

Table 6 displays the color coordinates of the breast and thigh meat of Pekin ducks fed various levels of dietary treatments. The first group samples (T1) indicated lower lightness (L^*) and high redness (a^*) ($P \leq 0.05$) with yellowness (b^*) of the breast, and females recorded lower lightness (L^*) and higher yellowness (b^*) than males because of the high presence of carotenoid pigment in the Alfalfa. On the other hand, the second group samples (T2) indicated lower lightness (L^*) and high redness (a^*) ($P \leq 0.05$) with yellowness (b^*) of the thigh, and males recorded higher in all traits than females. However, there were no significant differences ($P > 0.05$) in different feed levels on the yellowness color of breast and thigh, and in sex on the color of breast and thigh samples among birds. This observation corroborates the report of He et al., (15), observed that the b^* of breast muscles increased when Alfalfa meal was added to the diet, compared to the control ($P < 0.05$). However, a^* and L^* of the breast and thigh muscles were not affected by the Alfalfa meal addition. Compared with control chicken, Alfalfa meal chickens showed an increasing trend of b^* in their thigh muscles. In contrast, Michalczyk et al., (24) noted significantly higher redness a^* values of breast muscles in females than in males, while Gornowicz et al., (12) did not observe any significant differences in the redness of breast muscles between males and females. Different findings were concluded by Kokoszynski et al., (19) who reported that males showed significantly lower redness ($P = 0.002$) of breast muscles in Muscovy duck. The darker color from the analyzed Muscovy ducks resulted

from a significantly higher content of hem pigments compared to the breast muscles of Pekin ducks. This was probably related to better blood supply to the muscles in older Muscovy ducks. A study by Banaszak et al., (4) on two groups of Cherry valley meat ducks fed 650g/kg soybean meal and 689.89g/kg yellow lupin in concentrates, showed significantly greater ($P<0.05$) lightness (L^*) and yellowness (b^*) in the breast muscle of group 2 compared to group 1, but did not significantly differ for the color of the leg muscles. The greater lightness (L^*) and yellowness (b^*) of muscles from ducks fed yellow lupin may indicate a higher concentration of carotenoids in lupin as well as a higher intramuscular fat content in muscles. Similarly, a^* parameter values between the two groups in this study could be interpreted as a similar red muscular fiber composition in duck muscles. The lighter meat has a higher L^* value. This parameter can also be related to the pH of meat. Zhuang and Savage (42) found that dark meat has a lower L^* value and a higher pH, while light meat has a higher L^* value at a lower pH. Furthermore, Kokoszynski et al., (20), reported that significant ($P<0.05$) differences were observed in the color parameters of breast muscles and in the redness of leg muscles between males and females within two genetic groups of ducks. However, the study by Boz et al., (6) on Turkish geese varieties, indicated that significant differences ($P<0.05$) were observed in the yellowness of breast and thigh meat colors in lightness and yellowness between males and females.

3.3. pH values

Table 7 displays the pH values of the breast and thigh meat of Pekin ducks' sex-wise. pH values of different treatments decreased from pH_{60} (one-hour post-slaughter) to pH_{24} (24-hour post-mortem) and significantly ($P\leq 0.05$) from pH_f (post-freezing) in the breast. The breast and thigh of birds fed different percentages of Alfalfa pellet ration had lower pH than those from the control group, and breast females had lower pH than males and with significant differences between males and females in pH_{24} (24-hour post-mortem) and pH_f (post freezing) but, in the thigh, males had lower pH than females. The post-mortem

conversion of muscle glycogen to lactic acid may have contributed to the pH drop. The only source of energy for the post-mortem muscles in a bird that has been exsanguinated and subsequently suffers from hypoxia is anaerobic glycolysis, which eventually becomes necessary for the muscle cells' remaining metabolic processes (30). Glycogen stores are thus reduced when they are converted to lactic acid, resulting in a decrease in pH. The pH of muscle is an important indicator of various quality traits in meat. Similar findings were concluded by He et al., (15), who showed no significant differences in pH_{45} minutes and pH_{24} hours between treatments using different quantities of Alfalfa meal 25g/kg, 50g/kg, and 75g/kg diet of breast and thigh in the broiler. However, Banaszak et al., (4), reported similar findings too for non-significant ($P>0.05$) in pH_{15} and pH_{24} between two groups of Cherry valley meat ducks fed soybean meal (group 1) and yellow lupin (group 2) in diet, a change of protein source in duck diet had no significant effect on pH traits ($P>0.05$). The decrease in pH between 15 min and 24h post-mortem indicates correct glycolysis in breast muscles and was associated with the accumulation of lactic acid. Different findings were concluded by Addini et al., (2), who indicated significant differences ($P<0.05$) between different levels of Alfalfa 5% and 10% with control in pH of hybrid duck meat. Alfalfa's crude fiber is one of the feeds that affects the pH value. Higher alfalfa supplementation can reduce the meat's pH. Bernacki et al., (21), reported that the pH_{24} of breast muscles had significant differences ($P<0.05$) between females in different genetic groups, pH_{24} post-mortem of leg muscles did not differ shown. The leg muscles showed higher pH values measured 24 hours post-mortem compared with the breast muscles. Furthermore, Kokoszynski et al., (20), reported that the sex of birds did not have a significant ($P>0.05$) effect on the acidity (pH_{24}).

3.4. Proximate composition

There were no significant differences in breast and thigh proximate analysis, except breast protein content showed significant differences among treatments (Table 8). The breast meat from the control birds (T0) had the highest

protein and fat content which was significantly different (P<0.05) in protein content than the first treatment (T1).

Table 6. Effect of treatments and sex on the breast and thigh color for Pekin ducks at age 49 days

Color	Meat muscle	Overall mean	Treatments			SEM	Sex		SEM	P-value	
			T0	T1	T2		M	F		Treatments	Sex
L* Lightness	Breast	37.07	36.40ab	34.19b	40.64a	0.83	37.44	36.71	1.04	0.023	0.675
a* Redness		18.38	17.70ab	20.48a	16.95b	0.58	18.39	18.36	0.68	0.069	0.985
b* Yellowness		9.29	8.24	10.78	8.84	0.71	8.32	10.26	0.70	0.322	0.182
L* Lightness	Thigh	42.66	50.61a	41.96ab	35.40b	1.96	42.87	42.45	2.45	0.027	0.919
a* Redness		13.74	10.69b	14.15ab	16.38a	0.76	14.28	13.21	0.93	0.030	0.500
b* Yellowness		9.49	7.90	9.32	11.25	0.86	10.99	7.99	0.81	0.254	0.078

Means followed by the different letters in the same raw are significantly different (P<0.05). SEM: Pooled Standard Error of the Mean

Table 7. Effect of treatments and sex on the pH of breast and thigh muscles for Pekin ducks at age 49 days

pH measures	Meat muscle	Overall mean	Treatments			SEM	Sex		SEM	P-value	
			T0	T1	T2		M	F		Treatments	Sex
pH ₆₀	Breast	6.26	6.37	6.18	6.23	0.06	6.34	6.17	0.05	0.203	0.099
pH ₂₄		6.09	6.20	6.02	6.05	0.08	6.22a	5.95b	0.05	0.246	0.049
pH _f		6.02	6.08a	6.03b	5.93c	0.06	6.12a	5.91b	0.03	0.012	0.002
pH ₆₀	Thigh	6.49	6.56	6.43	6.47	0.02	6.47	6.50	0.03	0.171	0.493
pH ₂₄		6.33	6.35	6.31	6.34	0.01	6.32	6.35	0.01	0.350	0.199
pH _f		6.25	6.26	6.26	6.24	0.01	6.25	6.26	0.01	0.694	0.789

Means followed by the different letters in the same raw are significantly different (P<0.05). SEM: Pooled Standard Error of the Mean. pH₆₀: pH one-hour post-slaughter. pH₂₄: pH 24-hour post-mortem. pH_f: pH post freezing

Table 8. Effect of treatments and sex on the breast and thigh chemical composition for Pekin ducks at age 49 days

Chemical composition	Meat muscle	Overall mean	Treatments			SEM	Sex		SEM	P-value	
			T0	T1	T2		M	F		Treatments	Sex
Protein%	Breast	21.19	22.00a	20.09b	21.47a	0.42	20.47b	21.90a	0.41	0.019	0.012
Fat%		8.10	9.42	7.25	7.64	0.76	8.04	8.16	0.81	0.661	0.956
Ash%		1.33	1.39	1.28	1.33	0.05	1.31	1.36	0.05	0.793	0.744
Moisture%		70.12	68.14	71.44	70.79	0.72	70.88	69.37	0.87	0.328	0.395
Protein%	Thigh	19.66	20.21	19.28	19.5	0.27	19.52	19.80	0.30	0.553	0.697
Fat%		8.24	8.39	9.90	6.41	0.74	7.53	8.94	0.89	0.358	0.447
Ash%		1.10	1.13	1.09	1.06	0.03	1.11	1.09	0.03	0.735	0.779
Moisture%		71.08	70.40	69.39	73.44	0.62	71.25	70.90	1.01	0.278	0.841

Means followed by the different letters in the same raw are significantly different (P<0.05). SEM: Pooled Standard Error of the Mean

Nevertheless, there was no significant difference in breast and thigh fat contents (P>0.05) among birds fed diets T0, T1, and T2. The lower fat content observed in the T1 in the breast, and T2 in the thigh birds as compared to the control group, and in the male breast and thigh lower protein and fat content than females which were in line with the reports were recorded, the present results suggest that the replacement of alfalfa pellet ration (50%) to the basal diet recorded lower fat% of thigh meat than others in Pekin ducks. The different finding recorded by Addini et al., (2), who reported that alfalfa supplementation in commercial feed significantly affected the

water content of hybrid duck meat (P<0.05). The highest water content was at the alfalfa supplementation of 10%, while the lowest water content is at the level of alfalfa supplementation of 5%, and no significant differences (P<0.05) on the crude protein. Based on the measured amino acid contents, these results indicated that alfalfa can be a suitable alternative for other feed materials that provide the necessary amount of several amino acids for poultry including corn meal (lysine and tryptophan), soybean meal (methionine, cysteine, lysine, and tryptophan) and fish meal (methionine and lysine) (34). He et al., (15), reported that the effects of dietary

alfalfa meal addition on meat quality parameters of broilers that the thigh muscle crude protein was lower in the alfalfa meal group than in the control group ($P<0.05$), while there were no significant differences in terms of breast muscle crude protein. No significant differences were found in the Ether Extract of the thigh and breast muscles. In addition, the study by Palupi et al., (28) concluded that 0.5% of propionic acid added to the diet might improve the carcass quality of the broiler. In contrast, Galal et al., (11), who referred that the male ducks had significantly higher water and protein but had significantly lower fat and ash content in breast and thigh muscles compared to female ones. Kokoszynski et al., (19), reported that the sex of ducks had a considerable ($P<0.05$) effect on basic chemical composition (water%, protein%, fat%) of the breast and leg meat. However, the study by Kokoszynski et al., (20) indicated that males contained significantly less fat in breast muscles and significantly more water and protein, and less ($P<0.05$) fat in the leg muscles than in females. Boz et al., (6), who referred that crude fat in breast meat was higher in male geese ($P<0.05$) than in females. The effects of sex of other meat characteristics (Protein%, Ash%, Dry matter%) were not significant.

3.5. Vitamins

Table 9. Effect of treatments and sex on fat-soluble vitamins (ppm) in the breast and thigh for Pekin ducks at age 49 days

fat-soluble vitamins	Sample	Overall mean	Treatments			SEM	Sex		SEM	P-value	
			T0	T1	T2		M	F		Treatments	Sex
A	Breast	21.19	18.95b	22.87a	21.75a	1.34	18.87b	23.51a	0.83	0.010	0.003
D		0.80	0.60	0.88	0.91	0.22	0.43b	1.16a	0.08	0.320	0.032
E		0.54	0.36b	0.74a	0.54ab	0.09	0.39b	0.70a	0.08	0.075	0.039
K	Thigh	0.34	0.19b	0.47a	0.36ab	0.06	0.23b	0.44a	0.06	0.040	0.023
A		47.70	44.40b	50.66a	48.05ab	2.77	42.93b	52.47a	1.31	0.048	0.007
D		2.12	1.76b	2.45a	2.16a	0.12	1.92b	2.31a	0.14	0.035	0.036
E		1.91	1.67b	2.17a	1.91ab	0.43	1.17b	2.66a	0.11	0.084	0.003
K		1.12	0.88c	1.39a	1.11b	0.08	0.99b	1.25a	0.11	0.008	0.010

Means followed by the different letters in the same row are significantly different ($P<0.05$). SEM: Pooled Standard Error of the Mean

3.5.1. Fat-soluble vitamins: The effect of the replacement of alfalfa pellets to the basal diet on fat-soluble vitamins in Pekin duck's breast and thigh meat is shown in Table 9. The first group samples (T1) indicated high content (ppm) of fat-soluble vitamins than the second group (T2) and control (T0) respectively. The replacement of the alfalfa pellet ration to the basal diet had a significant influence ($P\leq 0.05$) on the fat-soluble vitamins except vitamin D which influenced not significantly ($P>0.05$) in comparison with control, and both breast and thigh females had a higher significantly ($P\leq 0.05$) content of fat-soluble vitamins than males (Table 9).

3.5.2. Water-soluble vitamins

The effect of the treatment and sex on water-soluble vitamins in the breast and thigh meat of Pekin duck is shown in Table 10. The second group samples (T2) indicated high content of water-soluble vitamins (ppm) than the first group (T1) and control (T0) respectively. The replacement of the alfalfa pellet ration to the basal diet had a significant influence ($P\leq 0.05$) on the water-soluble vitamin content in the breast and thigh in comparison with control, and both breast and thigh males had a higher significant content ($P\leq 0.05$) of most water-soluble vitamins than females (Table 10).

Table 10. Effect of treatments and sex on water-soluble vitamins (ppm) in the breast and thigh for Pekin ducks at age 49 days

Water-soluble vitamins	Sample	Overall mean	Treatments			SEM	Sex		SEM	P-value	
			T0	T1	T2		M	F		Treatments	Sex
B2	Breast	7.65	6.80b	7.95a	8.20a	0.20	8.00a	7.30b	0.31	0.005	0.007
B3		8.22	7.45b	8.30ab	8.90a	0.73	9.47a	6.97b	0.31	0.058	0.007
B5		4.58	3.85c	4.80b	5.10a	0.41	5.30a	3.87b	0.27	0.002	0.001
B6		6.48	5.75b	6.70a	7.00a	0.26	6.93a	6.03b	0.27	0.017	0.012
B12		3.55	2.90c	3.70b	4.05a	0.18	3.87a	3.23b	0.24	0.002	0.003
C	Thigh	6.50	5.75b	6.70ab	7.05a	0.34	7.07a	5.93b	0.28	0.054	0.026
B2		6.38	5.70c	6.55b	6.90a	0.18	6.70a	6.07b	0.25	0.002	0.003
B3		6.35	5.80b	6.50a	6.75a	0.13	6.57a	6.13b	0.20	0.024	0.039
B5		3.18	2.65c	3.30b	3.60a	0.11	3.37a	3.00b	0.20	0.004	0.008
B6		4.95	4.40b	5.10a	5.35a	0.13	5.17a	4.73b	0.20	0.024	0.039
B12		2.67	2.10b	2.80ab	3.10a	0.22	3.03	2.30	0.22	0.081	0.053
C		4.83	4.25b	4.95a	5.30a	0.10	5.00	4.67	0.22	0.020	0.063

Means followed by the different letters in the same row are significantly different ($P < 0.05$). SEM: Pooled Standard Error of the Mean

3.6. Cholesterol: Cholesterol content (mg/100gm) in the breast and thigh of the Pekin duck was shown in Table 11. The first group samples (T1) indicated lower content in the breast (34.50mg/100gm) of cholesterol than the second group (T2) (35.30mg/100gm) and control (T0) (45.55mg/100gm) respectively, while thigh Cholesterol content was lower in the first group (86.85mg/100gm) and the second group (T2) (87.75mg/100gm) than the control group (T0) (97.30mg/100gm). The replacement of the alfalfa pellet ration with the basal diet did not significantly influence ($P > 0.05$) cholesterol, and both the breast and thighs of males had a higher content of cholesterol than females (Table 11). In general, cholesterol content in the thigh was higher than breast which led to giving more beneficial health of the breast to humans than the thigh, especially for humans who had high blood pressure and cardiovascular diseases. Hadi et al., (13), reported that the addition of Perilla seeds which contain a high percentage of fiber, fat, and protein had a significant effect ($P < 0.05$) on cholesterol levels in duck meat. The addition of Perilla seed tends to increase cholesterol levels in duck meat. However, Banaszak et al., (4), reported that the using of lupin seeds as a replacement for soybean meal in the diet didn't have a significant effect on the cholesterol content in the breast muscle of Cherry Valley ducks. The study by Haraf et al., (14), indicated that the cholesterol content in the examined muscles ranged from 57.88 to 72.31mg/100g of goose muscles. Higher cholesterol content in thigh meat was observed by Boz et al., (6), for local Turkish goose varieties (74.95-77.85mg/100g). According to USDA (37), cholesterol content

in raw goose thigh should equal 84mg/100g of muscle. In publications by other authors, the cholesterol content in the femoral muscles of broiler chickens was different (60.5-68.15mg/100g of muscle) (8) or higher (156.3-192.2mg/100g of muscle) (31).

Table 11. Effect of treatments and sex on cholesterol (mg/100gm) in the breast and thigh muscles for Pekin ducks at age 49 days

Factors	Breast	Thigh
Overall mean	38.45	90.63
Treatments	NS	NS
T0	45.55±1.95	97.30±1.90
T1	34.50±5.00	86.85±6.75
T2	35.30±1.40	87.75±1.95
Sex	NS	NS
M	41.23±3.24	92.87±3.89
F	35.67±4.17	88.40±4.46

NS: Non-Significant ($p > 0.05$)

3.7. Fatty acid profile: The fatty acid profile of the breast and thigh is shown in Tables 12 and 13. The percentage of fatty acids was different among treatments. The content of saturated fatty acid in the breast and thigh (undecanoic, Myristic, Palmitic, and Stearic acid) was numerically and significantly ($P < 0.05$) higher in the second group (T2) than first (T1) and control group (T0) (Table 13). The highest content of saturated fatty acid in the breast and thigh was recorded in the T2 treatment (50% alfalfa pellet ration). While the lower content of saturated fatty acid in the breast and thigh in the group were fed only a basal diet and they were significantly different ($P \leq 0.05$). In our results, birds feed 50% alfalfa pellet ration (T2) showed the highest ($P \leq 0.05$) content of unsaturated fatty acid (oleic%, Linoleic%, Arachidonic%, Linolenic%,

Ericosenoic%) when compared to control group (T0). However, T1 and T2 did not show any significant differences. For the polyunsaturated fatty acids (PUFA) (Linoleic acid, Arachidonic, Linolenic), and monounsaturated fatty acids (MUFA) (Oleic, Ericosenoic), the control group recorded the lowest content of PUFA and MUFA, and the highest proportion was recorded for the second group (T2). The summation of PUFA, MUFA, SFA, n-3, n-6, and n-9 were varying in groups. The highest significant ($P \leq 0.05$) content of total PUFA, total MUFA, and total SFA in the breast and thigh, were recorded in the second group (T2). Also, breast and thigh females have higher numerically and significant ($P \leq 0.05$) content of most of all fatty acids than males (Table 12, 13). Saturated fatty acid (SFA) can have a negative impact on consumer healthy while monounsaturated fatty acid (MUFA) and polyunsaturated fatty acid (PUFA) are considered to have a positive effect on health. A similar finding in results by Wang et al., (38) on China Micro-ducks at 57 days of age, who used different levels of Alfalfa meal in diet (0, 65, 130, and 205g/kg), indicated that the proportion of fatty acids increased with increasing levels of dietary crude fibers. However, the study by Kwiecien et al., (22) on broilers that used two different doses of alfalfa protein concentrate 15g and 30g/kg, indicated that the chickens fed with alfalfa protein concentrate supplemented diets had higher contents ($P \leq 0.05$) of eicosadienoic and arachidonic acids and a lower level of saturated margaric (C17:0) and arachidic (C20:0) acids in the breast. However, had higher ($P < 0.05$) of eicosadienoic acids and arachidonic acids with a lower level ($P < 0.05$) of saturated palmitic and margaric acids as well as significantly lower content ($P < 0.01$) of arachidic acid in the thigh muscle, compared with the control group. Similarly, Uhlirva et al., (36), found that the effect of sex detected significant differences in the linoleic ($P < 0.001$) and α -linolenic acid ($p = 0.038$), total n-6 ($p = 0.016$), and total PUFA ($p = 0.028$)

contents, all of which were higher in females than males. Furthermore, the higher content of α -linolenic acid in females could be explained by the role of steroid hormones (29).

3.8. Amino acid profile: The percentage of amino acids (essential and non-essential) in the breast and thigh of Pekin ducks are presented in Tables 14 and 15. The essential amino acids of those treatments fed alfalfa pellet ration were slightly higher compared to the groups fed on basal diets. Methionine and Lysin content (%) in the breast of the first group (T1) and the second group (T2) was (4.03% 3.39%, T1 and 6.65%, 5.10%, T2) (Table 14), and in thigh content (3.40%, 2.31% T1 and 5.35%, 4.25%, T2) respectively (Table 15). The contents of threonine, histidine, and cysteine that are related to the meat flavor (41) were found to some extent differences among all groups. Significant differences were found between treatments in breast samples of Histidine, Phenylalanine, and Leucine (Table 14), and in thigh samples of Methionine, Histidine, Phenylalanine, and Leucine (Table 15) which the second group (T2) recorded higher significantly ($P \leq 0.05$) content than other groups. In addition, all essential amino acids in the breast and thigh have been recorded including Valine, Leucine, Phenylalanine, Histidine, Tyrosine, and Methionine, and their percentage compared to the basal diet groups (T0) was higher (Table 14, 15). Each protein in the meat is distinguished by its content of amino acids and its amount in meat affects flavor substances. Also, amino acids in the meat protein are the precursors of meat smell (41). Generally, breast and thigh samples of females recorded higher content of all amino acids than males. The study by Wang et al., (38), indicated that the proportion of some amino acids (Valine, Leucine and Isoleucine) increased significantly ($P < 0.05$) with increasing levels of dietary crude fibers compared with control. Boz et al., (6), noted that the effect of sex didn't have a significant ($P > 0.05$) effect on amino acids in the breast and thigh of Turkish goose varieties.

Table 12. Effect of treatments and sex on fatty acid profile (%) in the breast muscles for Pekin ducks at age 49 days

Means followed by the different letters in the same raw are significantly different (P<0.05). SEM: Pooled

Fatty acid	Treatments			SEM	Sex		SEM	P-value	
	T0	T1	T2		M	F		Treatments	Sex
undecanoic acid (C11:0)	0.64c	0.78b	0.92a	0.05	0.69b	0.87a	0.06	0.006	0.004
Myristic (C14:0)	0.32b	0.39b	0.60a	0.03	0.38	0.49	0.06	0.027	0.068
Palmitic (C16:0)	9.60c	10.60b	11.70a	0.41	9.93b	11.33a	0.43	0.009	0.007
Stearic (C18:0)	2.30c	3.10b	3.90a	0.21	2.73b	3.47a	0.33	0.005	0.008
Oleic (C18:1) ω ₉	10.55c	11.80b	13.65a	0.60	10.97b	13.03a	0.64	0.017	0.013
Linoleic (C18:2) ω ₆	6.15b	7.15b	8.85a	0.06	6.80b	7.97a	0.08	0.023	0.040
Linolenic (C18:3) ω ₃	0.23b	0.33ab	0.50a	0.35	0.30	0.40	0.56	0.071	0.142
Ericosenoic (C20:1) ω ₉	0.46b	0.57b	0.84a	0.03	0.52b	0.72a	0.06	0.041	0.049
Arachidonic (C20:4) ω ₆	0.29b	0.42ab	0.58a	0.03	0.38	0.48	0.06	0.032	0.082
∑ MUFA	11.01b	12.37b	14.49a	0.67	11.48b	13.75a	0.72	0.019	0.015
∑ PUFA	6.67c	7.90b	9.93a	0.40	7.48b	8.85a	0.68	0.009	0.017
∑ UFA	17.67c	20.26b	24.41a	1.05	18.96b	22.60a	1.39	0.001	0.001
∑ SFA	12.86b	14.87b	17.12a	0.70	13.74b	16.16a	0.87	0.003	0.003
∑ n-3	0.23b	0.33ab	0.50a	0.03	0.30	0.40	0.06	0.071	0.142
∑ n-6	6.44b	7.57b	9.43a	0.38	7.18b	8.45a	0.62	0.016	0.030
∑ n-9	11.01b	12.37b	14.49a	0.67	11.48b	13.75a	0.72	0.019	0.015

Standard Error of the Mean. ∑ MUFA: Total of monounsaturated fatty acids. ∑ PUFA: Total of polyunsaturated fatty acids. ∑ UFA: Total of unsaturated fatty acids. ∑ SFA: Total of saturated fatty acids. ∑ n-3: Total of omega 3. ∑ n-6: Total of omega 6. ∑ n-9: Total of omega 9

Table 13. Effect of treatments and sex on fatty acid profile (%) in the thigh muscles for Pekin ducks at age 49 days

Fatty acid	treatments			SEM	Sex		SEM	P-value	
	T0	T1	T2		M	F		Treatments	Sex
undecanoic acid (C11:0)	1.15c	1.65b	2.20a	0.14	1.43b	1.90a	0.22	0.021	0.034
Myristic (C14:0)	0.78b	0.93b	1.26a	0.04	0.91b	1.06a	0.10	0.013	0.047
Palmitic (C16:0)	16.95b	18.15b	20.70a	0.91	17.03b	20.17a	0.79	0.014	0.007
Stearic (C18:0)	6.40b	7.45ab	9.25a	0.72	6.50b	8.90a	0.61	0.071	0.035
Oleic (C18:1) ω ₉	19.05b	20.35b	22.85a	0.96	19.10b	22.40a	0.80	0.018	0.009
Linoleic (C18:2) ω ₆	14.45b	15.95b	18.95a	0.20	14.90b	18.00a	0.25	0.039	0.028
Linolenic (C18:3) ω ₃	0.42b	0.49b	0.77a	0.92	0.49b	0.62a	0.95	0.015	0.044
Ericosenoic (C20:1) ω ₉	0.88	1.17	1.95	0.04	1.04	1.63	0.08	0.146	0.156
Arachidonic (C20:4) ω ₆	0.58b	0.76b	1.12a	0.08	0.68b	0.96a	0.11	0.017	0.021
∑ MUFA	19.93c	21.52b	24.80a	1.12	20.14b	24.03a	1.01	0.0003	0.0001
∑ PUFA	15.45b	17.20b	20.84a	1.03	16.07b	19.58a	1.13	0.021	0.017
∑ UFA	35.37c	38.72b	45.64a	2.14	36.21b	43.60a	2.14	0.007	0.004
∑ SFA	25.28c	28.18b	33.41a	1.78	25.88b	32.03a	1.68	0.002	0.001
∑ n-3	0.42b	0.49b	0.77a	0.04	0.49b	0.62a	0.08	0.015	0.044
∑ n-6	15.03b	16.71b	20.07a	1.00	15.58b	18.96a	1.06	0.027	0.021
∑ n-9	19.93c	21.52b	24.80a	1.12	20.14b	24.03a	1.01	0.0003	0.0001

Means followed by the different letters in the same raw are significantly different (P<0.05). SEM: Pooled Standard Error of the Mean. ∑ MUFA: Total of monounsaturated fatty acids. ∑ PUFA: Total of polyunsaturated fatty acids. ∑ UFA: Total of unsaturated fatty acids. ∑ SFA: Total of saturated fatty acids. ∑ n-3: Total of omega 3. ∑ n-6: Total of omega 6. ∑ n-9: Total of omega 9

Table 14. Effect of treatments and sex on amino acid profile (%) in the breast muscles for Pekin ducks at age 49 days

Amino acid	Treatments			SEM	Sex		SEM	P-value	
	T0	T1	T2		M	F		Treatments	Sex
Essential amino acids									
Arginine (Arg)	3.13	4.05	5.62	0.34	3.81	4.72	0.55	0.120	0.231
Histidine (His)	2.19c	3.19b	4.98a	0.17	3.19	3.72	0.58	0.013	0.105
Leucine (Leu)	2.63b	3.54b	5.94a	0.29	3.57	4.50	0.70	0.019	0.076
Lysine (Lys)	3.25b	4.03b	6.65a	0.26	4.23	5.06	0.73	0.014	0.075
Methionine (Met)	2.46b	3.39b	5.10a	0.26	3.25	4.05	0.56	0.033	0.107
Phenylalanine (Phe)	2.80b	3.63b	6.11a	0.23	3.80	4.56	0.71	0.009	0.055
Valine (Val)	3.35	3.60	5.63	0.48	3.53	4.85	0.57	0.194	0.201
Non-Essential amino acids									
Alanine (Ala)	3.80	4.42	5.97	0.40	4.20	5.26	0.51	0.191	0.232
Aspartic (Asp)	2.84b	3.65ab	5.22a	0.30	3.50	4.30	0.52	0.098	0.225
Asparagine (Asn)	2.16b	2.85ab	4.91a	0.28	2.90	3.71	0.60	0.054	0.178
Glutamic (Glu)	2.63b	3.45ab	5.06a	0.30	3.30	4.12	0.53	0.085	0.201
Proline (Pro)	2.74	3.61	5.25	0.33	3.42	4.30	0.55	0.110	0.229
Serine (Ser)	2.89	3.38	5.10	0.34	3.31	4.27	0.51	0.122	0.194
Tyrosine (Tyr)	2.57b	3.32ab	5.37a	0.33	3.27	4.23	0.61	0.065	0.159

Means followed by the different letters in the same row are significantly different ($P < 0.05$). SEM: Pooled Standard Error of the Mean

Table 15. Effect of treatments and sex on amino acid profile (%) in the thigh muscles for Pekin ducks at age 49 days

Amino acid	Treatments			SEM	Sex		SEM	P-value	
	T0	T1	T2		M	F		Treatments	Sex
Essential amino acids									
Arginine (Arg)	2.55	2.78	4.83	0.33	2.93	3.83	0.54	0.099	0.202
Histidine (His)	1.61b	2.20b	4.12a	0.21	2.30	2.99	0.54	0.015	0.065
Leucine (Leu)	2.14b	2.67b	4.74a	0.22	2.83	3.52	0.57	0.017	0.082
Lysine (Lys)	2.80b	3.40b	5.35a	0.20	3.53	4.17	0.55	0.017	0.089
Methionine (Met)	1.81b	2.31b	4.25a	0.18	2.50	3.08	0.53	0.015	0.088
Phenylalanine (Phe)	2.27b	2.61b	5.01a	0.21	2.96	3.63	0.61	0.012	0.074
Valine (Val)	2.58	2.41	4.63	0.45	2.57	3.84	0.56	0.178	0.196
Non-Essential amino acids									
Alanine (Ala)	3.19	3.59	5.17	0.39	3.44	4.51	0.47	0.184	0.203
Aspartic (Asp)	1.95b	2.85ab	4.40a	0.24	2.73	3.40	0.52	0.062	0.211
Asparagine (Asn)	1.52b	1.79b	3.78a	0.21	2.05	2.68	0.51	0.029	0.124
Glutamic (Glu)	1.98b	2.77b	4.18a	0.19	2.69	3.26	0.46	0.026	0.116
Proline (Pro)	2.04b	2.72b	4.48a	0.24	2.72	3.44	0.52	0.038	0.127
Serine (Ser)	2.34	2.45	4.23	0.30	2.57	3.43	0.46	0.104	0.177
Tyrosine (Tyr)	1.97b	2.50b	4.46a	0.27	2.56	3.39	0.54	0.036	0.105

Means followed by the different letters in the same row are significantly different ($P < 0.05$). SEM: Pooled Standard Error of the Mean

Conclusion

It was concluded that the birds fed diet replacement with Alfalfa pellet ration beneficially influenced meat quality in comparison with those fed only basal diets. So, the replacement of the Alfalfa pellet ration influenced positively on most Physico-chemical traits. Feeding ducks with a 25% of Alfalfa pellet ration improved in terms of color breast, and fat-soluble vitamins with Cholesterol in both breast and thigh muscle while feeding ducks with a 50% Alfalfa pellet

ration improved most Physico-chemical traits. Furthermore, the breast and thighs of females had a higher content of protein%, fat%, amino acid profile, fatty acid profile, and fat-soluble vitamins than males.

REFERENCES

1. Abdulla, N. R., A. N. Mohd Zamri, A. B. Sabow, K. Y. Kareem, S. Nurhazirah, F. H. Ling, A. Q. Sazili, and T. C. Loh, 2017. Physico-chemical properties of breast muscle in broiler chickens fed probiotics, antibiotics or antibiotic-probiotic mix. Journal of Applied

- Animal Research, 45(1), p.64-70.
<https://doi.org/10.1080/09712119.2015.1124330>
- 2.Addini, S.A., B. Suwignyo, and C. Hanim, 2020. Supplementation Alfalfa (*Medicago sativa* L.) in commercial feed on physic and chemical quality meat of hybrid duck. In E3S Web of Conferences (Vol. 200, p. 03012). EDP Sciences.
<https://doi.org/10.1051/e3sconf/202020003012>
- 3.AOAC. 2000. Official methods of analysis of AOAC International (17th ed.) Gaithersburg (MD): AOAC. <https://scholar.google.com>
- 4.Banaszak, M., Kuźniacka, J., Biesek, J., Maiorano, G. and Adamski, M., 2020. Meat quality traits and fatty acid composition of breast muscles from ducks fed with yellow lupin. *Animal*, 14(9), p.1969-1975.
<https://doi.org/10.1017/S1751731120000610>
- 5.Biesiada-Drzazga B., 2012. Ducks. In: Jankowski J, editor. Poultry breeding and utilization. Warsaw, Poland: PWRiL; p. 377-396. (Cited by Kokoszyński et al., 2020
<https://doi.org/10.5194/aab-63-137-2020>)
- 6.Boz, M.A., F. Oz, U. S. Yamak, M. Sarica, and E. Cilavdaroglu, 2019. The carcass traits, carcass nutrient composition, amino acid, fatty acid, and cholesterol contents of local Turkish goose varieties reared in an extensive production system. *Poultry science*, 98(7), p.3067-3080.
<https://doi.org/10.3382/ps/pez125>
- 7.Dahl-Lassen, R., van Hecke, J., Jørgensen, H., Bukh, C., Andersen, B. and Schjoerring, J.K., 2018. High-throughput analysis of amino acids in plant materials by single quadrupole mass spectrometry. *Plant Methods*, 14(8), p.1-9.
<https://doi.org/10.1186/s13007-018-0277-8>
- 8.De Oliveira, J., S.V. Avanço, M. Garcia-Neto, and E.H.G. Ponsano, 2016. Composition of broilers meat. *Journal of Applied Poultry Research*, 25(2), p.173-181.
<https://doi.org/10.3382/japr/pfv095>
- 9.Duncan, D. B., 1955. Multiple Range Test and Multiple F Tests. *Biometrics* .11: 1-42.
<https://doi.org/10.2307/3001478>
- 10.FAOSTAT: Livestock primary, production quantity, duck meat. 2018. (accessed 2019 March 25).
<http://www.fao.org/faostat/en/data/QL>.
- 11.Galal, A., W.A.H. Ali, A.M.H. Ahmed, and K. A. Ali, 2011. Performance and carcass characteristics of Dumyati, Muscovy, Peking and Sudani duck breeds. *Egyptian Journal of Animal Production*, 48(2), p.191-202.
<https://dx.doi.org/10.21608/ejap.2011.94072>
- 12.Gornowicz, E. and L. Lewko, 2015. Ducks—Meat Production—Consumption—Tradition. National Research Institute of Animal Production: Kraków, Poland, p.1-44. (Cited by Kokoszyński et al., 2019
<https://doi.org/10.3390/ani9110963>)
- 13.Hadi, R.F., S. Sudiyono, S. N. Jannah, and W. Indriyani, 2019, December. The effect of enriched perilla seed (*Perilla frutescens* L.) in the diets on percentage of carcass and non-carcass, chemical quality, and levels of α -linoleic acid (ALA) of meat ducks. In AIP Conference Proceedings (Vol. 2199, No. 1). AIP Publishing.
<https://doi.org/10.1063/1.5141306>
- 14.Haraf Gabriela, J. Wołoszyn, A.Okruszek, Z. Goluch, M.Werenska, and M. Teleszko, 2021. The protein and fat quality of thigh muscles from Polish goose varieties. *Poultry Science*; 100:100992, 1-9.
<https://doi.org/10.1016/j.psj.2021.01.015>
- 15.He, G., Zhao, L., M.S.R. Shishir, Y. Yang, Q. Li, L. Cheng, and A. Guo, 2021. Influence of alfalfa meal, as a source of dietary fibre, on growth performance, development, pH of gastrointestinal tract, blood biochemical profile, and meat quality of broilers. *Journal of Applied Animal Research*, 49(1), pp.431-439.
<https://doi.org/10.1080/09712119.2021.2000417>
- 16.Kijowski J, 2001. Nutritional value and technological usefulness of meat in waterfowl. (in Polish). IV Intern. Targi „Farma Świń i Drobiu” Maj 18-20, Poznan, 2001.
<https://scholar.google.com>
- 17.Kokoszyński D, R Wasilewski, K Stęczny, Z Bernacki, K Kaczmarek, M Saleh, PD Wasilewski, and M. Biegniewska, 2015. Comparison of growth performance and meat traits in Pekin ducks from different genotypes. *Europ Poult Sci.*; 110:1–11.
<http://dx.doi.org/10.1399/eps.2015.110>
- 18.Kokoszyński, D., R. Wasilewski, K. Stęczny, M. Kotowicz, C. Hrnčar, and Arpašová, H., 2019. Carcass composition and selected meat quality traits of Pekin ducks

- from genetic resources flocks. *Poultry science*, 98(7), p.3029-3039.
<https://doi.org/10.3382/ps/pez073>
- 19.Kokoszyński, D., A. Wilkanowska, M. Saleh, M. Fik, and B. Bigorowski, 2021. Comparison of some meat and liver quality traits in Muscovy and Pekin ducks. *Journal of Applied Animal Research*, 49(1), p.118-124.
<https://doi.org/10.1080/09712119.2021.1895173>
- 20.Kokoszyński, D., H. Arpášová, C. Hrnčar, J. Žochowska-Kujawska, M. Kotowicz, and M. Sobczak, 2020. Carcass characteristics, chemical composition, physicochemical properties, texture, and microstructure of meat from spent Pekin ducks. *Poultry science*, 99(2), p.1232-1240.
<https://doi.org/10.1016/j.psj.2019.09.003>
- 21.Bernacki, Z., P. D. Wasilewski, J. Żochowska- Kujawska, M. Saleh, K. Stęczyński, M. Kmiecik, M. Kotowicz, R. Zwierzyński, D. Kokoszyński, M. Sobczak, and T. Bucek, 2020. Carcass, physicochemical and sensory characteristics of meat from genetic reserve ducks after two reproductive seasons. *South African Journal of Animal Science*, 50(1), p.55-68.
<https://hdl.handle.net/10520/EJC-1f6912ec67>
- 22.Kwiecień, M., A. Winiarska-Mieczan, A. Danek-Majewska, K. Kwiatkowska, and R. Krusiński, 2021. Effects of dietary alfalfa protein concentrate on lipid metabolism and antioxidative status of serum and composition and fatty acid profile and antioxidative status and dietetic value of muscles in broilers. *Poultry science*, 100(4), p.100974.
<https://doi.org/10.1016/j.psj.2020.12.071>
- 23.Maiorano, G., S. Knaga, A. Witkowski, D. Cianciullo, and M. Bednarczyk, 2011. Cholesterol content and intramuscular collagen properties of pectoralis superficialis muscle of quail from different genetic groups. *Poultry Science*, 90(7), p.1620-1626.
<https://doi.org/10.3382/ps.2010-01190>
- 24.Michalczuk, M., K. Damaziak, D. Pietrzak, A. Marzec, M. Chmiel, L. Adamczak, and T. Florowski, 2016. Influence of Housing System on Selected Quality Characteristics of Duck Meat. Chapter 1. Pekin duck. *Ann. Warsaw Univ. Life Sci.*; 55:89-97.
<https://bibliotekanauki.pl/articles/2912>
- 25.Maloney, S. K., and D. A. Gray . 1998. Characteristics of the febrile response in Pekin ducks. *Journal of Comparative Physiology B*, 168, 177-182.
- 26.NRC, 1994. *Nutrient Requirements of Poultry*, 9th rev. ed. Washington, DC: Natl. Acad. Press vol 74.
<https://books.google.iq/books>
- 27.Onbaşilar, E. E. and S. Yalcin, 2018. Fattening performance and meat quality of Pekin ducks under different rearing systems. *World's Poultry Science Journal*, 74(1), p.61-68.
<https://doi.org/10.1017/S004393391700099X>
- 28.Palupi, R., F. N. L. Lubis, and A. Suryani, 2022. Addition of propionic acid on nutrient digestibility and its effect on production and carcass quality of broiler. *Iraqi Journal of Agricultural Sciences*, 53(2):453-464.
<https://doi.org/10.36103/ijas.v53i2.1553>
- 29.Poureslami, R., K. Raes, G. M. Turchini, G. Huyghebaert, and S. De Smet, 2010. Effect of diet, sex and age on fatty acid metabolism in broiler chickens: n-3 and n-6 PUFA. *British Journal of Nutrition*, 104(2), pp.189-197.
<https://doi.org/10.1017/S0007114510000395>
- 30.Sabow, A.B., A.Q. Sazili, I. Zulkifli, Y. M. Goh, M.Z.A.A. Kadir, and K. D. Adeyemi, 2015. Physico-chemical characteristics of Longissimus lumborum muscle in goats subjected to halal slaughter and anesthesia (halothane) pre-slaughter. *Animal Science Journal*, 86(12), p.981-991.
<https://doi.org/10.1111/asj.12385>
- 31.Salma, U., A. G. Miah, T. Maki, M. Nishimura, and H. Tsujii, 2007. Effect of dietary *Rhodobacter capsulatus* on cholesterol concentration and fatty acid composition in broiler meat. *Poultry Science*, 86(9), p.1920-1926.
<https://doi.org/10.1093/ps/86.9.1920>
- 32.SAS, *Statistical Analysis System*, 2005. *User's Guide for Personal Computer*. Release 8.2 SAS Institute Inc. Cary, NC, USA.
<https://books.google.iq/books>
- 33.Seal, T. and K. Chaudhuri, 2017. High performance liquid chromatography method for the estimation of water soluble vitamin in five wild edible fruits consumed by the tribal people of north-eastern region in India. *International Journal of Current Microbiology and Applied Sciences*, 6(10), p.2900-2913.

<https://doi.org/10.20546/ijcmas.2017.610.343>
34. Suwignyo B., E. Suryanto, H. Sasongko, Y. Erwanto, and E. A. Rini, 2020. The Effect of fresh and hay alfalfa (*Medicago sativa* L.) supplementation on carcass quality of hybrid duck. IOP conference series: Earth and Environmental Science 478 (IOP Publishing):12024.
<https://doi.org/10.1088/17551315/478/1/012024>
35. Shafer, D., X. Chen, M. Sifri, M. Lilburn, D. Karcher, P. Cherry, P. Wakenell, S. Fraley, M. Turk, and G. S. Fraley. 2021. Centennial Review: History and husbandry recommendations for raising Pekin ducks in research or commercial production. Poultry Science, 100(8), 101241.
36. Uhlířová, L., E. Tůmová, D. Chodová, Z. Volek, and V. Machander, 2019. Fatty acid composition of goose meat depending on genotype and sex. Asian-Australasian journal of animal sciences, 32(1), p.137-143.
<https://doi.org/10.5713/ajas.17.0672>
37. USDA. 2019. U. S. Department of Agriculture, Agricultural Research Service. Food Data Central. Accessed Mar. 2020. fdc.nal.usda.gov.
<https://farmdocdaily.illinois.edu/2020/04/review-wing-usda-funding-in-the-cares-act.html>
38. Wang, S.P., D.M. Luo, G.W. Liu, M. J. Yin, C. F. Zhou, X. L. Zhao, and Y. B. Guo, 2019. Responses of chemical composition,

amino acid and fatty acid profiles of breast muscle to dietary crude fibre levels in China Micro-Ducks. South African Journal of Animal Science, 49(3), p.485-493.
<https://doi.org/10.4314/sajas.v49i3.9>
39. Xue, X., J. You, and P. He, 2008. Simultaneous determination of five fat-soluble vitamins in feed by high-performance liquid chromatography following solid-phase extraction. Journal of chromatographic science, 46(4), p.345-350.
<https://doi.org/10.1093/chromsci/46.4.345>
40. Zhang, H., Z. Wang, and O. Liu, , 2015. Development and validation of a GC-FID method for quantitative analysis of oleic acid and related fatty acids. Journal of Pharmaceutical Analysis, 5(4), p.223-230.
<https://doi.org/10.1016/j.jpha.2015.01.005>
41. Zhao Y., C. Li, L. Zhu, S. Zhan, H. Wang, J. Wei, , et al., 2011. Clinical Diagnosis and Pathogenic Identification of Egg Drop-Death Syndrome in Duck. J. Yangzhou Univ. Agric. Life Sci. Ed. 32, 11–14.
<https://www.cabidigitallibrary.org/doi/full/10.5555/20113375023>
42. Zhuang, H. and E. M. Savage, 2010. Comparisons of sensory descriptive flavor and texture profiles of cooked broiler breast fillets categorized by raw meat color lightness values. Poultry science, 89(5), p.1049-1055.
<https://doi.org/10.3382/ps.2009-00422>