

USING *Rhabditis blumi* SUDHAUS AS BIOLOGICAL AGENTS TO CONTROL THE PALP BORER, ARABIAN RHINOCEROS BEETLE, *Oryctes agamemnon arabicus*

A. M. Tariq

Assis. Prof.

Instit. Medic. Techn. Al-Mansur, Middle Technical University, Baghdad, Iraq

ahmadtarek2001@yahoo.com

ABSTRACT

This study was aimed to use pathogenicity of entomopathogenic nematodes (EPN) *Rhabditis blumi* Sudhaus (Nematoda: Rhabditida) against palm borer Arabian Rhinoceros Beetle (ARB), *Oryctes agamemnon arabicus* was evaluated in the lab. And date palm orchards during 2015 - 2017. In the lab tests, EPN was used against the larvae and adults as a direct spray and as treated food (pieces of fresh tissue of the frond bases) at a rate of 0, 500, 1000, 1500 Infective Juveniles (IJs) per mL of *R. blumi*. Results indicated that EPN caused 89%, 61% and 25%, 20% mortality when used as a direct spray and as treated food on larvae and adults of ARB, respectively. Results of field experiments showed that injection of 50 mL per palm tree with a concentration of 1500 IJs/mL of *R. blumi* inflicted about 45.5% mortality in ARB larvae infesting the tree. Meanwhile, the Population density of ARB larvae reduced to 45.8%, 59.6% during the first and second year of treated date palm trees by injection method respectively. The results of this investigation illustrate the possibility of using EPN *R. blumi* as a biocontrol agent for managing borers in date palm orchards under field conditions.

Keywords: biocontrol, palm borers, *Oryctes agamemnon arabicus*, entomopathogenic nematode, *Rhabditis blumi*

طارق

مجلة العلوم الزراعية العراقية - 2020: 51(2): 657-664

استخدام *Rhabditis blumi* Sudhaus (Nematoda:Rhabditida) كعامل مقاومة احيائي لمكافحة خنفساء وحيدة

القرن العربية *Oryctes Agamemnon arabicus*

احمد محمد طارق

استاذ مساعد

المعهد التقني الطبي /المنصور, الجامعة التقنية الوسطى, بغداد, العراق .

المستخلص

الحفارات من الآفات الخطيرة على النخيل في العراق والبلدان الأخرى التي يزرع فيها نخيل التمر تسبب هذه الآفات ضررا بالغاً في أشجار النخيل وتقلل انتاجها كما ونوعاً وتسبب ضعف في الجذع مما يؤدي الى موت النخلة وانكساره. تم تقييم فاعلية النيما تودا الممرضة للحشرات *Rhabditis blumi* مختبرياً وحقلياً كعامل مكافحة احيائية في مكافحة خنفساء وحيدة القرن العربية *Oryctes agamemnon arabicus* خلال 2015 – 2017 , وقد استعملت النيما تودا ضد البالغات واليرقات بطريقتي الرش المباشر ومعاملة الغذاء (قطع طرية من كرب النخيل) , وقد استعملت النيما تودا بالتركيز 0 , 500 , 1000 , 1500 طور معدي (IJs) في كل مل. اشارت نتائج الفاعلية تحت ظروف المختبر بأن النيما تودا أعطت نسبة قتل 89% , 61% , 25% , 20% عندما استعملت بالرش المباشر وفي معاملة الغذاء على اليرقات والبالغات على التوالي . اما تجارب الفاعلية الحقيقية فقد أعطت فاعلية قتل في اليرقات 45.5% عندما استعملت بتركيز 1500 طور معدي (IJs) لكل مل. كما اشارت النتائج حصول انخفاض في الكثافة السكانية ليرقات خنفساء وحيدة القرن العربية بنسبة 45.5% , 59.6% وذلك خلال السنة الأولى والثانية بعد معاملة النخيل على التوالي .من النتائج المتحصل عليها يمكن ان تخدم في استعمال النيما تودا *Rhabditis blumi* كعامل مكافحة احيائية في برنامج إدارة حفارات النخيل في بساتين نخيل التمر عند الظروف الحقلية.

كلمات مفتاحية: المقاومة الحيوية, حفار النخيل, *Oryctes agamemnon arabicus*, النيما تودا الممرضة للحشرات, *Rhabditis blumi*

*Received:22/7/2019, Accepted:8/10/2019

INTRODUCTION

Palm borers, especially *Oryctes* spp. are considered as an economically important insect pest of date palm trees in Iraq and most adapted to climatic conditions of the region (12 and 4). Arabian Rhinoceros Beetle (ARB), *Oryctes agamemnon arabicus* causes severe damages to the bases of fronds and bunches making long tunnels inside the tissue, which are acting as weakening and breaking factors for these parts (13). Many control methods have been used through different application methods: spray, drench and injection (5, 88, and 11) against to date palm pests. Khalaf (14) Reported that the application of thiamethoxam and imidacloprid against ARB larvae resulted in 85.8%, 100% mortality in injection method compared with 75%, 80% in the drench method respectively. Khudair (15) Tested locally isolated entomopathogenic fungi *Metarhizium anisopliae* and *Beauveria bassiana* against ARB larvae and reported a high mortality rate among larvae reaching up to 100% after 29 days of treatment under lab conditions. Entomopathogenic nematodes will only be widely used as pest control products when they become available on demand by the different clients, commercial growers and small farmers (16). Rhabditid nematodes are generally recognized as bacterivores and often associated with invertebrates, their relationship with invertebrates is known as necromancy, which means waiting for the death host (6, 20, 23, 28 and 26). Khalaf (14) Tested entomopathogenic nematodes (EPN), *Rhabditis blumi*, and the entomopathogenic fungi (EPF), *Beauveria bassiana* as biocontrol agents against larvae and adults of ARB, *O. agamemnon arabicus*. Biological control potentials of *R. blumi* against 5 coleopteran species and 5 lepidopteran species was evaluated by Park (19). This study aimed to investigate the efficacy of EPN, *Rhabditis blumi* as entomopathogenic biocontrol agents against palm borers, *Oryctes* spp. especially ARB, *O. agamemnon arabicus* under laboratory and field conditions. Another objective was to investigate the feasibility of application and persistence in date palm orchards.

MATERIALS AND METHODS

Borers and nematodes cultures: Samples of ARB *O. agamemnon arabicus* were obtained

from the lab. Colony reared on natural foods (palm tissues, fresh frond bases pieces) at $25\pm 2^{\circ}\text{C}$, complete darkness and 65% relative humidity in the IPM Center, Directorate of Agricultural Research, and Ministry of Science and Technology (14). *R. blumi* was obtained from the lab. Colony reared on larvae of wax moth *Galleria mellonella* in labs of IPM center (14).

Laboratory experimental treatments

The laboratory trials were conducted in Biological Control Dept. of IPM Center. Laboratory experiments included using four concentrations (0, 500, 1000 and 1500 IJs per mL) of *R. blumi* as direct spray on larvae and adults of ARB or mixed with their food (pieces of frond bases tissue). Fresh food pieces were added regularly through the period of treatment to keep enough fresh food to the larvae. Five replicates, 6 larvae/rep as larval treatment and four replicates, 5 adults/rep as adult's treatment were used for each treatment as for laboratory experiments. Larval and adults mortalities were counted in all treatments after 24, 72, 96 and 120 hours of treatment.

Field experimental treatments

Field experiments, injection of 50 mL solution of EPN 1500 IJs/mL was done through tree trunk using 50 mL syringes after drilling holes with a brad point drill-bit (diameter, 20mm and length, 200mm) 1m above the ground level (Fig. 1-A, B). The same number of trees were left as control treatment in each orchard. Five replicates (trees) were used for each treatment. Larvae of ARB in trees crown were collected after four weeks of injection EPN, dead and live larvae were counted in each treatment and kept in plastic containers (Fig. 1-C). Also, fresh frond bases tissues were collected and healthy larvae of *G. mellonella* were added to test if it contains EPN, *R. blumi* in each treatment. Samples of tissue were taken after 4 weeks of treatment for examination and to explore the presence and movement of EPN through plant tissue. Larval mortalities that infected by EPN in each treatment were counted.

Efficacy and persistent of entomopathogenic nematode *Rhabditis blumi* in date palm orchards: Efficacy and persistence of EPN *R. blumi* were studied in

two date palm orchards, the first one was untreated (control) orchard, the second orchard was treated with EPN mixed in 50 mL at a concentration of 1500 IJs by used injection methods in ten tree trunk. EPN distribution and efficacy in reduction of ARB Larvae population density were counted by calculating larvae in ten palm tree Brem variety in each orchard before treatment and during the first and second year after treatment.

Experimental design and data analysis

The experimental designs implemented were complete randomized design and randomized complete block design with four replicates . Genstat program was implied in statistical analysis and determine the significances efficacies. Henderson-Tilton, s formula (1955) (10) was used to calculate corrected mortality

efficacies% on larval and adults ARB treated with EPN in lab treatments, while Schneider-Orelli, s formula (1981) (10) was applied for measuring efficacies of the EPN tested on ARB at field treatment.

RESULTS AND DESCUSSION

Results in Table 1 indicates that the mortality percentage of ARB larvae after direct spray under laboratory conditions with EPN *R. blumi* revealed that highest concentration (1500 IJs/mL) caused the highest percentage of mortality 89% after 120 hr., while the lower concentrations of 1000 and 500 IJs/mL recorded mortality of 79% and 57% respectively. Meanwhile, mortalities reached 50%, 54% and 61% at concentrations of 500, 1000 and 1500 IJs/mL respectively in experiments when EPN was mixed with larval food (table1).

Table 1. Effect of entomopathogenic nematode *Rhabditis blumi* as a biocontrol agent against palm borer, *Oryctes agamemnon arabicus* larvae under laboratory conditions

Treatment method	Concentration of <i>R. blumi</i> (IJs/mL)	Total of larvae treated	%Corrected mortality(accumulation) After(hr)											
			24			72			96			120		
			Larva dead	% mortality	% Efficacy	Larva dead	% mortality	% Efficacy	Larva dead	% mortality	% Efficacy	Larva dead	% mortality	% Efficacy
Direct spray	Control (water)	30	0	---	---	0	---	---	1	---	---	2	---	---
	500	30	13	43	43	17	57	57	17	57	55	18	60	57
	1000	30	21	70	70	23	77	77	23	77	76	24	80	79
	1500	30	25	83	83	26	87	87	27	90	89	27	90	89
Treated diet (frond bases)	Control (water)	30	0	---	---	0	---	---	1	---	---	1	---	---
	500	30	15	50	50	15	50	50	16	53	52	16	53	50
	1000	30	14	47	47	15	50	50	17	57	55	17	57	54
	1500	30	16	53	53	16	53	53	19	63	62	19	63	61

Adult's mortality for direct spray and treated food were 0%, 15%, 20%, 25% and 0%, 10%, 15%, 20% at concentration 0, 500, 1000 and 1500 IJs per mL respectively (Table 2).

Table 2. Effect of entomopathogenic nematode *Rhabditis blumi* as a biocontrol agent against palm borer, *Oryctes agamemnon arabicus* adults under laboratory conditions.

Treatment method	Concentration of <i>R. blumi</i> (IJs/mL)	Total of adults treated	%Corrected mortality(accumulation) After(hr)											
			24			72			96			120		
			adult dead	% mortality	% Efficacy	adult dead	% mortality	% Efficacy	adult dead	% mortality	% Efficacy	adult dead	% mortality	% Efficacy
Direct spray	Control (water)	20	0	---	---	0	---	---	0	---	---	0	---	---
	500	20	0	0	0	0	0	0	3	15	15	3	15	15
	1000	20	1	5	5	3	15	15	4	20	20	4	20	20
	1500	20	3	15	15	5	25	25	5	25	25	5	25	25
Treated diet (frond bases)	Control (water)	20	0	---	---	0	---	---	0	---	---	0	---	---
	500	20	0	0	0	2	10	10	2	10	10	2	10	10
	1000	20	2	10	10	2	10	10	3	15	15	3	15	15
	1500	20	3	15	15	3	15	15	4	20	20	4	20	20

Field efficacy results indicated that the mortality percentage of ARB larvae after trunk injection with 50 mL in concentration 1500 IJs

per mL reached 45.5% (calculated in the crown tree only) after 4 weeks of treatment (Table 3).

Table.3 Effect of entomopathogenic nematode, *Rhabditis blumi* as biocontrol agents against on palm borer, *Oryctes agamemnon arabicus* larvae under field conditions

Treatment (Trunk injection)	Number of larvae per five trees (in tree crown only)			
	Before treatment	After 4 weeks of treatment		%corrected mortality
		Life	Dead	
Control	Unknown	35	0	---
50 mL (1500 IJs per mL) per tree	Unknown	12	10 (dead and new place of larva)	45.5

The use of EPN *R. blumi* in date palm orchards as a biocontrol agent against *Oryctes* larvae caused a reduction in population density

reached 45.8% and 59.6% after one month and during the second year of treatment respectively (Table 4).

Table 4. Distribution and efficacy of entomopathogenic nematode *Rhabditis blumi* in date palm orchards in Alsweera in Middle of Iraq

Treatment (orchard)	Number of larvae per ten tree in crown tree only (Brem variety)				
	Before treatment	During the first year after treatment		During the second year after treatment	
		Life larva	Life larva	% Reduction in population	Life larva
Control Orchard No 1	120	131	---	115	---
Without Nematode Orchard No.2	93	55	45.8	36	59.6
Nematode 1500 IJs					

Results of field studies revealed that there was an acceptable efficacy of applying local isolate of EPN *R. blumi* (isolated locally from Iraqi date palm orchards ecosystem) as biocontrol agents. The EPN *R. blumi* can persist habitat causing more reduction in the population density of ARB, *O. agamemnon arabicus* larvae. In addition, results indicated that the EPN, *R. blumi* solution could translocate through date palm tissue after injection in the trunk. Park (19) Reported that EPN *R. blumi* against major cruciferous insect pests and evaluated pathogenicity in lab and greenhouse and showed that EPN caused high mortality rate in larvae. Entomopathogenic nematodes have certain advantages over chemical as control agents; it's a non-polluting agent and thus environmentally safe and acceptable (9). There are many reports for *rhabditid* nematodes causing mortality of various invertebrates' species: beetles, termite, millipede and rice yellow stem borer (1, 7, 22,

23, and 25). (2, 3,11,24 and 27) they found that the Infective Juveniles of *rhabditid* nematodes enter an invertebrate, remain until it is dead, and complete their development by feeding on bacteria growing inside the cadaver of insect, and it usually possesses some attributes of a potential biological control agents, such as short life cycle, easy culture condition with bacteria, high fecundity and virulence, and good association with invertebrate pathogenic bacteria. EPN *Rhabditis blumi* showed significant mortality against Arabian rhinoceros beetle *Oryctes agamemnon arabicus*, but the mortality rate of larvae was higher than that of adults in lab trails at direct spray or treated food. When EPN was mixed with 50 ml of water at a concentration of 1500 IJs/ml and injected in the trunk, a moderate mortality rate was reported among ARB larvae. However, dispersal and efficacy were increased in treated orchards after one year of treatment.




Figure 1. (A)- Inspection site for ARB larvae, (B) - location of EPN Injection site, (C)- Inspection of palm tree crown for collecting ARB larvae.

REFERENCES

1. Abood, S.M., M.R. Abood and A.A. Jasim. 2018. Manufacturing and testing of date palm vibration motorized frauds cutter. Iraqi Journal of Agricultural Sciences –1028:49(3):763- 73 1
2. Abood, I.D. and H.S.H. Yasien .2016. Detection resistance genes of root knot nematode Mi in some of pure lines of tomato indeterminate growth. Abood, I.D. Iraqi Journal of Agricultural Sciences – 47(5):1321-1327
3. Al- sahaf, F.H. , Abed Y.M. , Saleh , F.F. , and H.S.M. Khierallah ,. 2017. Influence of spraying by some plant growth regulators on date palm salt tolerance enhancement... The Iraqi Journal of Agricultural Sciences – 48(1): 236-241,2017
4. Al_shali, M.A.M, Al_kalyani, A.I.N, Dayinna and K.Yardi 2019 Can Industrialization affect heavy metals bio concentration in date palm tree farms in the Sultanate of Oman. Iraq Journal of Agricultural Sciences. 2019:50 (Special Issue) : 152-172
5. Al-Jboory, I. J., A. I. Al.Sammariya, J. F. Whaib and W. A. Ahmed, 2001. Evaluation of thiamethoxam in a different application technique to control Dubas bug *Ommatissus lybicus*. J. Arab Pl. Prot., 19:47- 53
6. Blaxter, M. L., P. De Ley, J. R. Gary, L. X. Liu, P. Scheldemman and A. Vierstraete. 1998. A molecular evolutionary framework for the phylum Nematoda. Nature, 392: 71-75
7. Carta, L. K. and J. C. Osbrink. 2005. *Rhabditis rainai* n. sp. (Nematoda: Rhabditida)

- associated with the Formosan subterranean termite, *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Nematology*, 7: 863-879
8. Felsot, S. A., 2002. Application of New Generation Systemic Insecticide Through Drip Irrigation Systems: Case Study With Imidacloprid. Research and Extension regional water quality conference, pp:3
9. Georgis, R. 1990. Formulation and Application Technology in Gaugler, R., H. K. Kaya, eds. *Entomopathogenic Nematodes in Biological Control*. Boca Raton. FL: CRC Press. , pp: 173- 191
10. Henderson, C. F. and E. W. Tilton. 1955. Tests with acaricides against the brow wheat mite, *J. Econ. Entomol* 48:157-161
11. [http:// www. researchgate. net/ publication/ 228425523](http://www.researchgate.net/publication/228425523). Accessed on 19 Jan. 2017.
- Timmeren, S.V., C.J. Wise and R. Isaacs, 2012. Soil application of neonicotinoid insecticide for control insect pests in wine grape, published online in Wiley online library. *Pest Mang. Sci.*, 68:537-542
12. Khalaf M. Z., A. K Shbar., M. H. Al-Seria, R. A. Sami, and F. H. Naher 2011. Some aspects of biology and control methods of fruit stalk borer *Oryctes elegans* Prell (Coleoptera: Scarabaeidae). *Journal of Agricultural Science and Technology a* 1:142-147
13. Khalaf M. Z., and A. A Al-Taweel,. 2015. Palm borers in Iraqi environment: species-damages- methods of control. *J. of the Blessed Tree*, 07:54-64
14. Khalaf, M. Z., H. F. Alrubaei, F. H. Naher and M. Dh. Jumaa. 2016. Biological control of the date palm tree borers, (Coleoptera: Scarabidae: Dynastinae. Book of Proceedings VII International Scientific Agriculture Symposium (Agrosym2016). Jahorina, Bosnia and Herzegovina, October 06-09, 1561-1566
15. Khudhair, M. W., M. Z. Khalaf, H. F. Alrubaei A. K. Shbar, B. S Hamad. and H. S. Khalaf 2015. Evaluating the virulence of *Metarhizium anisopliae* (Deuteromycotina: Hyphomycetes) and *Beauveria bassiana* (Ascomycota: Hypocreales) isolates to Arabian rhinoceros beetle, *Oryctes agamemnon arabicus*. *J. of Entomological and Acarological Research* 47: 117-122
16. Mwaniki, S. W., J. H. Nderitu, F. Olubayo and J. W. kimenju. 2013. Mass production of entomopathogenic nematodes using Silkworm, *Bombyx mori* L. for management of key agricultural pests. 12th KARI Scientific Conference Proceedings 2010: 759-763
17. Padmakumari, A. P., J. S. Prasas, G. Katti and M. Sankar. 2007. *Rhabditis* sp. (*Oscheius* sp.), abiotic agent against rice yellow stem borer, *Scirpophaga incertulas*. *Inian J. Plant protection*. 35: 2-28
18. Park, H. W., H. H. Kim, S.H. Youn, T. S. Shin, A. L. A. L. Bilgrami, M. R. Cho and C. S. Shin. 2012. Biological control potentials of insect-parasitic nematode *Rhabditis blumi* (Nematoda: Rhabditida) for major cruciferous vegetable insect pests. *Applied Entomol. Zool*. 47: 389-397
19. Park, H. W., Kim, M. R. Cho, T. J. Kang, S. J. Ahu, S. W. Jeon, and A. L. Bilgrami. 2012. Evaluation of biological potentials of *Rhabditis blumi* (Nematoda: Rhabditida) against 10 insect species. *Journal Information service System* 37:235-239
20. Park, H. W., Y. O. Kim, J. Ha, S. H. Youn, H. H. Kim, A. L. Bilgrami and C. S. Shin. 2011. Effects of associated bacteria on the pathogenicity and reproduction of the insect parasitic nematode *Rhabditis blumi* (Nematoda: Rhabditida). *Can. J. Microbiol* 57:750-758
21. Püntener, W. 1981. Manual for Field Trials in Plant Protection 2nd. ed. Agricultural Division, Ciba-Geigy Limited ,pp.37
22. Richter, S. 1993. Phoretic association between between the dauer juveniles of *Rhabditis stammeri* and life history stages of the burying beetle *Nicrophorus vespilloides* (Coleoptera: Silphidae). *Nematologica*, 39: 346 – 355
23. Schulte, F 1989. The association between *Rhabditis nectomena* Sudhaus and Schutle, 1989 (Nematoda: Rhabditidae) and native and introduced millipedes in South Australia. *Nematologica*, 35: 82-89
24. Shblawy, L. M. and Al-Jorany R.S. 2018. Sunn pest species and its hibernation sites in Diyala Governorate / IRAQ. *Iraqi Journal of Agricultural Sciences* –49(6):0002-0088
25. Smart, G. C. and K. B. Nguyen. 1994. *Rhabditis Pheropsophi* (Rhabditida: Rhabditidae). *J. Nematology*. 26: 19-24

26. Stock, S. P., A. M. Caiceda and P. A. Calatayud. 2005. *Rhabditis (Oscheius) Colombiana* (Nematoda: Rhabditidae), a necromenic associate of the subterranean burrower bug *Cyrtomenus bergi* (Hemiptera: Cydnidae) from Cauca Valley, Colombia. *Nematology*, 7: 417-373
27. Sudhaus, W. and F. Schulte. 1989. *Rhabditis necromena* (Nematoda:Rhabditidae) from South Australian diplopoda with notes on its sibling *Rhabditis myriophila* poinar, 1986 and *Rhabditis caulleryi* Manupas. *Nematologica*, 35: 15-24
28. Wilson, M. J., D. M. Glen, S. K. George, and R. C. Butler. 1993. Mass cultivation and storage of Rhabditid Nematode *Phasmarhabditis hermaphrodita*, a biocontrol agent for slige. *Biocontrol Sci. Technol.* 3: 513-521.